



RESEARCH ARTICLE

Forbidden induced subgraphs for graphs and signed graphs with eigenvalues bounded from below

Zilin Jiang ¹ and Alexandr Polyanskii ²

¹School of Mathematical and Statistical Sciences, and School of Computing and Augmented Intelligence, Arizona State University, Tempe, AZ 85281, USA; E-mail: zilinj@asu.edu (Corresponding author)

²Department of Mathematics, Emory University, Atlanta, GA 30322, USA; E-mail: apolian@emory.edu

Received: 6 February 2022; **Revised:** 14 April 2025; **Accepted:** 2 September 2025

2020 Mathematical Subject Classification: *Primary* – 05C50; *Secondary* – 05D10, 52C35

Abstract

The smallest eigenvalue of a graph is the smallest eigenvalue of its adjacency matrix. We show that the family of graphs with smallest eigenvalue at least $-\lambda$ can be defined by a finite set of forbidden induced subgraphs if and only if $\lambda < \lambda^*$, where $\lambda^* = \rho^{1/2} + \rho^{-1/2} \approx 2.01980$, and ρ is the unique real root of $x^3 = x + 1$. This resolves a question raised by Bussemaker and Neumaier. As a byproduct, we find all the limit points of smallest eigenvalues of graphs, supplementing Hoffman's work on those limit points in $[-2, \infty)$.

We also prove that the same conclusion about forbidden subgraph characterization holds for signed graphs. Our impetus for the study of signed graphs is to determine the maximum cardinality of a spherical two-distance set with two fixed angles (one acute and one obtuse) in high dimensions. Denote by $N_{\alpha,\beta}(d)$ the maximum number of unit vectors in $\mathbb{R}^d$ where all pairwise inner products lie in $\{\alpha, \beta\}$ with $-1 \leq \beta < 0 \leq \alpha < 1$. Very recently Jiang, Tidor, Yao, Zhang, and Zhao determined the limit of $N_{\alpha,\beta}(d)/d$ as $d \rightarrow \infty$ when $\alpha + 2\beta < 0$ or $(1 - \alpha)/(\alpha - \beta) \in \{1, \sqrt{2}, \sqrt{3}\}$, and they proposed a conjecture on the limit in terms of eigenvalue multiplicities of signed graphs. We establish their conjecture whenever $(1 - \alpha)/(\alpha - \beta) < \lambda^*$.

Contents

1	Introduction	2
2	Forbidden subgraphs for $\mathcal{G}(\lambda)$	5
2.1	Proof of Theorem 1.3 for $\lambda < 2$	5
2.2	Proof of Theorem 1.3 for $\lambda \in [2, \lambda^*)$	9
2.3	Proof of Theorem 1.3 for $\lambda \geq \lambda^*$	13
3	Forbidden subgraphs for $\mathcal{G}^\pm(\lambda)$	19
3.1	Proof of Theorem 1.5 for $\lambda < 2$	20
3.2	Proof of Theorem 1.5 for $\lambda \in [2, \lambda^*)$	23
3.3	Proof of Theorem 1.5 for $\lambda \geq \lambda^*$	25
4	Spherical two-distance sets	25
5	Concluding remarks	32
A	Computer-assisted proofs	33
	References	35

1. Introduction

A fundamental problem in spectral graph theory is the classification and characterization of graphs with bounded eigenvalues. When we talk about eigenvalues of a graph we always refer to its adjacency matrix. In this paper, we study the families of graphs with eigenvalues bounded from below. Let $\mathcal{G}(\lambda)$ be the family of graphs with smallest eigenvalue at least $-\lambda$. For the sake of comparison, we mention the family $\mathcal{G}'(\lambda)$ of graphs with spectral radius (or, equivalently, largest eigenvalue) at most λ .

Remark. Since we rarely work with subgraphs that are not induced, all subgraphs are induced throughout this paper. We refer to subgraphs that are not necessarily induced as *general subgraphs*.

The Cauchy interlacing theorem implies that both $\mathcal{G}'(\lambda)$ and $\mathcal{G}(\lambda)$ are closed under taking subgraphs. It is a natural question to ask whether it is possible to define each of these families by a finite set of forbidden subgraphs.

Definition 1.1. Given a family $\mathcal{G}$ of graphs that is closed under taking subgraphs, a family $\mathcal{F}$ of graphs is a *forbidden subgraph characterization* of $\mathcal{G}$ if the family $\mathcal{G}$ consists exactly of graphs that do not contain any member of $\mathcal{F}$ as a subgraph, and a graph F is a *minimal forbidden subgraph* for $\mathcal{G}$ if F itself is not in $\mathcal{G}$ but every proper subgraph of F is in $\mathcal{G}$.

Note that the most economical forbidden subgraph characterization of $\mathcal{G}$ consists precisely of the minimal forbidden subgraphs for $\mathcal{G}$. Thus the existence of a finite forbidden subgraph characterization of $\mathcal{G}$ is equivalent to the finiteness of the minimal forbidden subgraphs for $\mathcal{G}$. In 1992 Bussemaker and Neumaier made the following remark in [6, p 599]:

It would be interesting to know the set of numbers $m, -m$ such that $\mathcal{G}_m^\#$ [the set of minimal forbidden subgraphs for $\mathcal{G}'(m)$] or $\mathcal{G}_{-m}^\#$ [the set of minimal forbidden subgraphs for $\mathcal{G}(m)$] are finite; however, these seem to be very difficult problems.

The specific families $\mathcal{G}'(2)$ and $\mathcal{G}(2)$ are well understood. One of the earliest results dates back to 1970 when Smith [29] determined all the connected graphs in $\mathcal{G}'(2)$ – they are general subgraphs of the extended Dynkin diagrams in Figure 1. The family $\mathcal{G}(2)$ is much richer and more complex – it contains not only all the graphs in $\mathcal{G}'(2)$, but also all the line graphs.¹ The classification of $\mathcal{G}(2)$ culminated in a beautiful theorem of Cameron, Goethals, Seidel, and Shult [7] who related $\mathcal{G}(2)$ to root systems that occur in the classification of semisimple Lie algebras (see Theorem 2.17). We refer the reader to the monograph [12] for a comprehensive account of $\mathcal{G}(2)$.

These classification theorems can be used to establish quantitative answers to Bussemaker and Neumaier's problems. For $\mathcal{G}'(2)$, Cvetković, Doob, and Gutman [9, Theorem 2.8] determined that there are 18 minimal forbidden subgraphs. For $\mathcal{G}(2)$, Rao, Singhi, and Vijayan [26, Theorem 4.1] observed that the number of vertices in any minimal forbidden subgraph is at most 37, which was eventually perfected to 10 by Kumar, Rao, and Singhi [23]. A computer search by Bussemaker and Neumaier [6, p 596] established that there are, in total, 1,812 minimal forbidden subgraphs for $\mathcal{G}(2)$.

In a recent work, the authors of the current paper resolved the first problem of Bussemaker and Neumaier.²

Theorem 1.2 (Theorem 1 of Jiang and Polyanskii [20]). *For every integer $m \geq 2$, let β_m be the largest root of $x^{m+1} = 1 + x + \cdots + x^{m-1}$, and let $\alpha_m := \beta_m^{1/2} + \beta_m^{-1/2}$. The family $\mathcal{G}'(\lambda)$ of graphs with*

¹A line graph $L(H)$ of a graph H is obtained by creating a vertex per edge in H , and connecting two vertices if and only if the corresponding edges in H have a vertex in common. The adjacency matrix of $L(H)$ can be written as $B^T B - 2I$, where B is the vertex-edge incidence matrix of H , hence the smallest eigenvalue of $L(H)$ is at least -2 .

²The original statement of [20, Theorem 1] determines the set of λ for which the subfamily of connected graphs in $\mathcal{G}'(\lambda)$ (or equivalently, the entire family $\mathcal{G}'(\lambda)$) can be defined by a finite set of forbidden *general* subgraphs. Using the well-known fact that the spectral radius of G_1 is at most that of G_2 whenever G_1 is a general subgraph of G_2 , one can show that if $\mathcal{G}$ is a forbidden general subgraph characterization for $\mathcal{G}'(\lambda)$, then the family $\{H : \exists G \in \mathcal{G} \text{ s.t. } G \text{ is a general subgraph of } H \text{ on the same vertex set}\}$ is a forbidden (induced) subgraph characterization for $\mathcal{G}'(\lambda)$, and so the original statement of [20, Theorem 1] is equivalent to Theorem 1.2.




Figure 1. Maximal connected graphs with spectral radius at most 2. The number of vertices is one more than the given index. In particular, $\tilde{D}_4$ is actually a star with four leaves.

spectral radius at most λ has a finite forbidden subgraph characterization if and only if $\lambda < \lambda'$ and $\lambda \notin \{\alpha_2, \alpha_3, \dots\}$, where

$$\lambda' := \lim_{m \rightarrow \infty} \alpha_m = \varphi^{1/2} + \varphi^{-1/2} = \sqrt{2 + \sqrt{5}} \approx 2.05817,$$

and φ is the golden ratio $(1 + \sqrt{5})/2$.

The first main theorem of this paper resolves the remaining problem of Bussemaker and Neumaier – $\mathcal{G}(\lambda)$ enjoys a straightforward threshold phenomenon.

Theorem 1.3. The family $\mathcal{G}(\lambda)$ of graphs with smallest eigenvalue at least $-\lambda$ has a finite forbidden subgraph characterization if and only if $\lambda < \lambda^*$, where

$$\lambda^* := \rho^{1/2} + \rho^{-1/2} \approx 2.01980,$$

and ρ is the unique real root of $x^3 = x + 1$.

Remark. The constants defined in Theorems 1.2 and 1.3 satisfy $2 < \lambda^* = \alpha_2 < \alpha_3 < \dots < \lambda'$. The constant ρ defined in Theorem 1.3, known as the *plastic number*, is the smallest Pisot–Vijayaraghavan number.³

As a byproduct, we determine all the limit points of the set of smallest eigenvalues of graphs. Let Λ_1 consist of $\lambda \in \mathbb{R}$ such that $-\lambda$ is the smallest eigenvalue of some graph. Here we invert the smallest eigenvalues of graphs similarly to how we define $\mathcal{G}(\lambda)$. Before our work, Hoffman [18] characterized all the limit points of Λ_1 in $(-\infty, 2]$. His result involves a technical qualification, which was conjectured to be always true, and was later established by Greaves et al. [15, Theorem 1]. Doob [14, Theorem 9] observed that every real number in $\{\alpha_2, \alpha_3, \alpha_4, \dots\} \cup [\lambda', \infty)$ is a limit point of Λ_1 , and conjectured that α_2 (which equals λ^*), $\alpha_3, \alpha_4, \dots$ are the only limit points of Λ_1 in $(2, \lambda')$. We refute this conjecture by finding all the limit points of Λ_1 in $(2, \lambda')$.

Corollary 1.4. For every $\lambda > 2$, the negative number $-\lambda$ is a limit point of the set of smallest eigenvalues of graphs if and only if $\lambda \geq \lambda^*$.

We next turn our attention to *signed graphs*, which are graphs whose edges are each labeled by $+$ or $-$. Throughout the paper we decorate variables for signed graphs with the $\pm$ superscript. When we talk about eigenvalues of a signed graph $G^\pm$ on n vertices, we refer to its *signed adjacency matrix* – the $n \times n$ matrix whose (i, j) -th entry is 1 if ij is a positive edge, -1 if ij is a negative edge, and 0 otherwise.

It still makes sense to speak of forbidden subgraph characterization of a family of signed graphs with eigenvalues bounded from below. Our second main theorem establishes the same threshold phenomenon for the family of signed graphs with smallest eigenvalue at least $-\lambda$.

³The exact value of ρ is $\sqrt[3]{(9 + \sqrt{69})/18} + \sqrt[3]{(9 - \sqrt{69})/18}$.

Theorem 1.5. *The family $\mathcal{G}^\pm(\lambda)$ of signed graphs with smallest eigenvalue at least $-\lambda$ has a finite forbidden subgraph characterization if and only if $\lambda < \lambda^*$.*

Notice that if $\mathcal{F}^\pm$ is a finite forbidden subgraph characterization of $\mathcal{G}^\pm(\lambda)$, then the set of all-positive signed graphs in $\mathcal{F}^\pm$ is a finite forbidden subgraph characterization of $\mathcal{G}(\lambda)$. Therefore, for $\lambda < \lambda^*$, Theorem 1.5 implies Theorem 1.3. However, in Section 2, we still provide the complete proof of Theorem 1.3 to illustrate the key ideas.

Finally, we turn our attention to the largest eigenvalue of a signed graph. We denote the eigenvalues of a signed graph $G^\pm$ by $\lambda_1(G^\pm) \leq \lambda_2(G^\pm) \leq \dots$ in ascending order, and by $\lambda^1(G^\pm) \geq \lambda^2(G^\pm) \geq \dots$ in descending order. Observing that for every signed graph $G^\pm$, $\lambda_1(G^\pm) \geq -\lambda$ if and only if $\lambda^1(-G^\pm) \leq \lambda$, where $-G^\pm$ reverses all the edge signs of $G^\pm$, we obtain an immediate consequence of Theorem 1.5.

Corollary 1.6. *The family $\mathcal{G}^\mp(\lambda)$ of signed graphs with largest eigenvalue at most λ has a finite forbidden subgraph characterization if and only if $\lambda < \lambda^*$.*

Our motivation to understand the forbidden subgraph characterization of $\mathcal{G}^\mp(\lambda)$ comes from the problem of determining the maximum size of a *spherical two-distance set* with two fixed angles (one acute and one obtuse) in high dimensions. For fixed $-1 \leq \beta < 0 \leq \alpha < 1$, let $N_{\alpha,\beta}(d)$ denote the maximum number of unit vectors in $\mathbb{R}^d$ where all pairwise inner products lie in $\{\alpha, \beta\}$.

In the special case $\alpha = -\beta$, which corresponds to *equiangular lines*, recent work [5, 2, 20] culminated in a solution [21] of Jiang, Tidor, Yao, Zhang and Zhao to the problem of determining $N_{\alpha,-\alpha}(d)$ for sufficiently large d . We refer the reader to [21, Section 1] for earlier developments on equiangular lines with fixed angles in high dimensions.

For the general case $-1 \leq \beta < 0 \leq \alpha < 1$, in their subsequent work [22], Jiang et al. proposed a conjecture on the limit of $N_{\alpha,\beta}(d)/d$ as $d \rightarrow \infty$. To state their conjecture, we need the following spectral graph theoretic quantity.

Definition 1.7. Given $\lambda > 0$ and $p \in \mathbb{N}$, define the quantity

$$k_p(\lambda) = \inf \left\{ \frac{|G^\pm|}{\text{mult}(\lambda, G^\pm)} : \chi(G^\pm) \leq p \text{ and } \lambda^1(G^\pm) = \lambda \right\},$$

where $|G^\pm|$ is the number of vertices of $G^\pm$, $\text{mult}(\lambda, G^\pm)$ is the multiplicity of λ as an eigenvalue of $G^\pm$, and $\chi(G^\pm)$ is the chromatic number of the signed graph $G^\pm$.

We postpone the definition of $\chi(G^\pm)$, which takes values in $\mathbb{N}^+ \cup \{\infty\}$, to Section 4 (see Definition 4.1). We now state the conjecture on $N_{\alpha,\beta}(d)$.

Conjecture 1.8 (Conjecture 1.11 of Jiang et al. [22]). *Fix $-1 \leq \beta < 0 \leq \alpha < 1$. Set $\lambda = (1 - \alpha)/(\alpha - \beta)$ and $p = \lfloor -\alpha/\beta \rfloor + 1$. Then*

$$N_{\alpha,\beta}(d) = \begin{cases} \frac{k_p(\lambda)d}{k_p(\lambda)-1} + o(d) & \text{if } k_p(\lambda) < \infty, \\ d + o(d) & \text{otherwise.} \end{cases}$$

Conjecture 1.8 was confirmed in [22] for $p \leq 2$ and for $\lambda \in \{1, \sqrt{2}, \sqrt{3}\}$ separately. Building on the framework developed there, for all $\lambda < \lambda^*$, we establish Conjecture 1.8 as an application of Corollary 1.6, and we reduce the error term $o(d)$ to a constant depending only on α and β .

The rest of the paper is organized as follows. We prove Theorem 1.3 and Corollary 1.4 in Section 2, and we prove Theorem 1.5 in Section 3. Part of the proofs are computer-assisted with validated numerics. In Appendix A we explain our computer-aided proof and how anyone can recreate it independently. In Section 4 we give an application of Theorem 1.5 to spherical two-distance sets. In Section 5, we discuss open problems related to the classification of graphs in $\mathcal{G}(\lambda^*) \setminus \mathcal{G}(2)$, and a possible way to establish more instances of Conjecture 1.8.




Figure 2. $E_{2,n}$.

2. Forbidden subgraphs for $\mathcal{G}(\lambda)$

We break the proof of Theorem 1.3 into three cases $\lambda < 2$, $\lambda \in [2, \lambda^*)$ and $\lambda \geq \lambda^*$, where $\lambda^* \approx 2.01980$ is defined as in Theorem 1.3. It is worth pointing out the spectral graph theoretic interpretation of the peculiar constant λ^* .

Proposition 2.1 [20, Lemma 5(e)]. *For every $n \in \mathbb{N}^+$, define the graph $E_{2,n}$ as in Figure 2. As $n \rightarrow \infty$, $\lambda^1(E_{2,n})$ increases to λ^* , or equivalently $\lambda_1(E_{2,n})$ decreases to $-\lambda^*$.⁴*


2.1. Proof of Theorem 1.3 for $\lambda < 2$

Hoffman demonstrated several sequences of graphs $G_1, G_2, \dots$ such that G_m is a subgraph of G_{m+1} for every m and $\lim_{m \rightarrow \infty} \lambda_1(G_m) \leq -2$. To state these results, we introduce the following notions.

Definition 2.2. Given a nonempty vertex subset A of a graph F , $\ell \in \mathbb{N}$, and $m \in \mathbb{N}^+$,

- (a) the *path extension* (F, A, ℓ) is obtained from F by adding a path $v_0 \dots v_\ell$ of length ℓ , and connecting v_0 to every vertex in A ;⁵
- (b) the *path-clique extension* (F, A, ℓ, K_m) is further obtained from (F, A, ℓ) by adding a clique of order m , and connecting every vertex in the clique to v_ℓ ;
- (c) the *clique extension* (F, A, K_m) is obtained from F by adding a clique of order m , and connecting every vertex in the clique to every vertex in A .

The following figure consists of schematic drawings of the path extension (F, A, ℓ) , the path-clique extension (F, A, ℓ, K_m) , and the clique extension (F, A, K_m) .



We compile some of Hoffman's computation [18] and two classical results in the following lemma.

Lemma 2.3. *Denote by C_n the cycle of length n , and $V_2(C_n)$ a set of two adjacent vertices of C_n . The path-clique extensions and clique-extensions of C_n satisfy:*

- (c1) $\lim_{m \rightarrow \infty} \lambda_1(C_n, V_2(C_n), \ell, K_m) \leq -2$ for fixed $n \geq 3$ and $\ell \in \mathbb{N}$;
- (c2) $\lim_{m \rightarrow \infty} \lambda_1(C_n, V_2(C_n), K_m) = -2$ for fixed $n \geq 3$.

Denote by K_n the complete graphs with n vertices. The path-clique extensions of K_n satisfy:

- (k) $\lim_{m \rightarrow \infty} \lambda_1(K_m, V(K_m), \ell, K_m) = -2$ for fixed $\ell \in \mathbb{N}$.

Denote by $\overline{K}_2$ the null graph with 2 vertices. The path-clique extensions and the clique extensions of $\overline{K}_2$ satisfy:

- (n1) $\lim_{m \rightarrow \infty} \lambda_1(\overline{K}_2, V(\overline{K}_2), \ell, K_m) = -2$ for fixed $\ell \in \mathbb{N}$;
- (n2) $\lim_{m \rightarrow \infty} \lambda_1(\overline{K}_2, V(\overline{K}_2), K_m) = -2$.

⁴The equivalence is due to the fact that the spectrum of a bipartite graph (e.g., a tree) is symmetric around 0.

⁵When $\ell = 0$, the path of length ℓ is simply a single vertex.




Figure 3. The claw graph C and the diamond graph D .

Denote by P_n the path of length n (with $n + 1$ vertices), and S_n the star with n leaves. Their smallest eigenvalues satisfy:

- (p) $\lim_{n \rightarrow \infty} \lambda_1(P_n) = -2$;
- (s) $\lambda_1(S_n) = -\sqrt{n}$.

Remark. In Lemma 2.3, (c1) to (n2) are taken directly from [18, Lemmas 2.4 to 2.9], and (p) and (s) follow from the classical results $\lambda^1(P_n) = 2 \cos(\pi/(n + 2))$ and $\lambda^1(S_n) = \sqrt{n}$.

Lemma 2.3 allows us to build a finite set of forbidden subgraphs for $\mathcal{G}(\lambda)$. To state the result, we introduce the following definition.

Definition 2.4 (Extension family). Given a graph F and $\ell, m \in \mathbb{N}^+$, the *extension family* $\mathcal{X}(F, \ell, m)$ of F consists of the path-extension (F, A, ℓ) , the path-clique extension (F, A, ℓ_0, K_m) , and the clique extension (F, A, K_m) , where A ranges over the nonempty vertex subsets of F , and ℓ_0 ranges over $\{0, \dots, \ell - 1\}$.

Lemma 2.5. For every $\lambda < 2$, there exist $\ell, m \in \mathbb{N}^+$ such that the extension families $\mathcal{X}(C, \ell, m)$ and $\mathcal{X}(D, \ell, m)$ are both disjoint from $\mathcal{G}(\lambda)$, where C is the claw graph and D is the diamond graph (see Figure 3).

Proof. From Lemma 2.3(p), we obtain $\ell \in \mathbb{N}^+$ such that $P_\ell \notin \mathcal{G}(\lambda)$. Clearly, for every nonempty A , the path extensions (C, A, ℓ) and (D, A, ℓ) , each of which contains P_ℓ as a subgraph, are not in $\mathcal{G}(\lambda)$. With hindsight, using Lemma 2.3(c1, c2, n1, n2), we choose $m \in \mathbb{N}^+$ such that none of the following graphs

$$(C_3, V_2(C_3), \ell_0, K_m), (C_3, V_2(C_3), K_m), (\overline{K}_2, V(\overline{K}_2), \ell_0, K_m), (\overline{K}_2, V(\overline{K}_2), K_m), \quad (1)$$

where $\ell_0 \in \{0, \dots, \ell + 1\}$, is in $\mathcal{G}(\lambda)$.

It suffices to prove that every clique extension and every path-clique extension in the extension families $\mathcal{X}(C, \ell, m)$ and $\mathcal{X}(D, \ell, m)$ contains one of the graphs in (1) as a subgraph. Label the vertices of C and D as in Figure 3, and pick an arbitrary $\ell_0 \in \{0, \dots, \ell - 1\}$. The clique extension (C, A, K_m) and the path-clique extension (C, A, ℓ_0, K_m) respectively contain

$$\begin{aligned} &(\overline{K}_2, V(\overline{K}_2), 0, K_m) \text{ and } (\overline{K}_2, V(\overline{K}_2), \ell_0 + 1, K_m) \text{ when } 0 \in A \text{ and } |A \cap \{1, 2, 3\}| \leq 1; \\ &(\overline{K}_2, V(\overline{K}_2), 1, K_m) \text{ and } (\overline{K}_2, V(\overline{K}_2), \ell_0 + 2, K_m) \text{ when } 0 \notin A \text{ and } |A \cap \{1, 2, 3\}| = 1; \\ &(\overline{K}_2, V(\overline{K}_2), K_m) \text{ and } (\overline{K}_2, V(\overline{K}_2), \ell_0, K_m) \text{ when } |A \cap \{1, 2, 3\}| \geq 2, \end{aligned}$$

and the clique extension (D, A, K_m) and the path-clique extension (D, A, ℓ_0, K_m) respectively contain

$$\begin{aligned} &(\overline{K}_2, V(\overline{K}_2), 0, K_m) \text{ and } (\overline{K}_2, V(\overline{K}_2), \ell_0 + 1, K_m) \text{ when } A \subseteq \{0, 2\} \text{ and } A \neq \emptyset; \\ &(C_3, V_2(C_3), 0, K_m) \text{ and } (C_3, V_2(C_3), \ell_0 + 1, K_m) \text{ when } A \in \{\{1\}, \{3\}\}; \\ &(\overline{K}_2, V(\overline{K}_2), K_m) \text{ and } (\overline{K}_2, V(\overline{K}_2), \ell_0, K_m) \text{ when } A \supseteq \{1, 3\}; \\ &(C_3, V_2(C_3), K_m) \text{ and } (C_3, V_2(C_3), \ell_0, K_m) \text{ when } |A \cap \{0, 2\}| \geq 1 \text{ and } |A \cap \{1, 3\}| = 1. \end{aligned}$$

Therefore the extension families $\mathcal{X}(C, \ell, m)$ and $\mathcal{X}(D, \ell, m)$ are both disjoint from $\mathcal{G}(\lambda)$. $\square$

The next result shows that forbidding a star and an extension family of F effectively forbids F itself in every sufficiently large connected graph.

Lemma 2.6. *For every graph F and $k, \ell, m \in \mathbb{N}^+$, there exists $N \in \mathbb{N}$ such that for every connected graph G with more than N vertices, if no member in $\{S_k\} \cup \mathcal{X}(F, \ell, m)$ is a subgraph of G , then neither is F .*

Proof. With hindsight, we choose $N = vd^\ell$, where d is the Ramsey number $R(k, 2^{v+1}m + v)$, and v is the number of vertices of F . Suppose that G is a connected graph with more than N vertices that contains no member in $\{S_k\} \cup \mathcal{X}(F, \ell, m)$ as a subgraph. Assume for the sake of contradiction that the subgraph of G induced on a vertex subset, say V , is isomorphic to F .

Since G is a connected graph with more than vd^ℓ vertices, we claim that either there exists a vertex u at distance at least ℓ from V , or the maximum degree of G is at least d . Indeed, assume for the sake of contradiction that every vertex of G is within distance $\ell - 1$ from V , and the maximum degree of G is less than d . For $i \in \{0, 1, \dots, \ell - 1\}$, let V_i be the set of vertices at distance i from V . Clearly $V_0 = V$, and so $|V_0| = v$. Using the assumption on the maximum degree of G , one can inductively show that $|V_i| \leq vd^i$. Therefore the number of vertices in G expressed as $\sum_{i=0}^{\ell-1} |V_i|$ is at most $\sum_{i=0}^{\ell-1} vd^i = v(d^\ell - 1)/(d - 1) \leq vd^\ell$, which yields a contradiction.

We break the rest of the proof into two cases.

Case 1: There exists a vertex u at distance at least ℓ from V . Thus the subgraph $G[V]$ and a shortest path from V to u contains the path extension (F, A, ℓ) as a subgraph for some nonempty $A \subseteq V$, which is a contradiction.

Case 2: The maximum degree of G is at least d . Let u be a vertex of G with the maximum degree, and let $N(u)$ be the set of neighbors of u in G . Since S_k is not a subgraph of G , the subgraph $G[N(u)]$ cannot have an independent set of size k . Because $d = R(k, 2^{v+1}m + v)$, the subgraph $G[N(u)]$ contains a clique of order $2^{v+1}m + v$, and so there exists a clique $G[U]$ of size $2^{v+1}m$ that is vertex-disjoint from F .

We further partition the vertices in U as follows. For every subset A of V , let $U(A)$ be the set of vertices in U that are adjacent to all vertices in A and not adjacent to any vertex in $V \setminus A$. By the pigeonhole principle, there exists a subset A of V such that $|U(A)| \geq 2m$. If A is nonempty, then $G[V \cup U(A)]$ contains the clique extension (F, A, K_m) as a subgraph, which is a contradiction.

Hereafter we may assume that $|U(\emptyset)| \geq 2m$, hence the distance between V and $U(\emptyset)$ is at least 2. Let $v_0 \dots v_{\ell_0}$ be a shortest path between V and $U(\emptyset)$, where v_0 and v_{ℓ_0} are respectively at distance 1 from V and $U(\emptyset)$. Let A_1 be the nonempty set of vertices in V that are adjacent to v_0 , and denote by v_{ℓ_0+1} an arbitrary vertex in $U(\emptyset)$ that is adjacent to v_{ℓ_0} . We may assume that $\ell_0 < \ell - 1$ because otherwise $G[V \cup \{v_0, \dots, v_{\ell_0}, v_{\ell_0+1}\}]$ would contain the path extension (F, A_1, ℓ) as a subgraph, which is a contradiction. Let A_2 be the nonempty set of vertices in $U(\emptyset)$ that are adjacent to v_{ℓ_0} . If $|A_2| \geq m$, then $G[V \cup \{v_0, \dots, v_{\ell_0}\} \cup A_2]$ contains the path-clique extension (F, A_1, ℓ_0, K_m) as a subgraph. Otherwise $|U(\emptyset) \setminus A_2| > m$, and so $G[V \cup \{v_0, \dots, v_{\ell_0}, v_{\ell_0+1}\} \cup (U(\emptyset) \setminus A_2)]$ contains the path-clique extension $(F, A_1, \ell_0 + 1, K_m)$. $\square$

Combining Lemmas 2.5 and 2.6, we obtain a finite set of forbidden subgraphs for $\mathcal{G}(\lambda)$ that forbids the claw graph and the diamond graph in every sufficiently large connected graph. The following result, which is an immediate consequence of [27, Theorem 4], gives a sufficient condition for line graphs.

Theorem 2.7 (Theorem 4 of van Rooij and Wilf [27]). *Every graph that contains neither the claw graph nor the diamond graph as a subgraph is a line graph.*

Furthermore, we can bootstrap to obtain a finite set of forbidden subgraphs for $\mathcal{G}(\lambda)$ that forces every sufficiently large connected graph to be the line graph of a tree whose complexity is uniformly bounded.

Lemma 2.8. *For every $\lambda < 2$, there exist $N \in \mathbb{N}$ and a finite family $\mathcal{F}_1$ that is disjoint from $\mathcal{G}(\lambda)$ such that for every connected graph G with more than N vertices, if G contains no member in $\mathcal{F}_1$ as a subgraph, then there exists a rooted tree $H \in \mathcal{T}_N$ such that G is the line graph of H , where $\mathcal{T}_N$ is the family of rooted trees such that every connected component obtained from removing the root has at most N vertices.*

Proof. We obtain $\ell, m \in \mathbb{N}^+$ from Lemmas 2.3 and 2.5 such that the following family

$$\begin{aligned}\mathcal{F}_1 := & \{S_4, P_\ell\} \cup \mathcal{X}(C, \ell, m) \cup \mathcal{X}(D, \ell, m) \\ & \cup \{(C_n, V_2(C_n), \ell_0, K_m) : n \in \{3, \dots, \ell + 1\}, \ell_0 \in \{0, \dots, \ell - 1\}\} \\ & \cup \{(C_n, V_2(C_n), K_m) : n \in \{3, \dots, \ell + 1\}\} \\ & \cup \{(K_m, V(K_m), \ell_0, K_m) : \ell_0 \in \{0, \dots, \ell - 1\}\}\end{aligned}$$

is disjoint from $\mathcal{G}(\lambda)$.

We obtain N_0 from Lemma 2.6 such that for every connected graph G with more than N_0 vertices, if no member in $\{S_4\} \cup \mathcal{X}(C, \ell, m) \cup \mathcal{X}(D, \ell, m)$ is a subgraph of G , then neither is C nor D . With hindsight, we choose $N = \max(N_0, (m + 1)^{\ell+2})$. Suppose that G is a connected graph with more than N vertices, and suppose that no member in $\mathcal{F}_1$ is a subgraph of G . By our choice of N_0 , we know that neither C nor D is a subgraph of G .

Theorem 2.7 implies that G is the line graph of another connected graph, say H . Since P_ℓ is not a subgraph of G , the path $P_{\ell+1}$ cannot be a general subgraph of H . In particular, the diameter⁶ of H is at most ℓ . Let d be the maximum degree of H . If $d < m + 2$, then H has at most $(m + 1)^{\ell+1}$ vertices, and at most $(m + 1)^{\ell+2}$ edges, and so G has at most $(m + 1)^{\ell+2}$ vertices, which is a contradiction.

We may assume that $d \geq m + 2$. Let u be a vertex of H with degree d . We claim that H is a tree. Suppose on the contrary that H contains a cycle as a general subgraph. Since H does not contain $P_{\ell+1}$ as a general subgraph, the length of any cycle in H cannot exceed $\ell + 1$. Suppose in addition that u is on a cycle in H . Let C_n be a shortest cycle containing u , where $n \in \{3, \dots, \ell + 1\}$. By the choice of C_n , the vertex u has at least $d - 2 \geq m$ neighbors outside C_n . Thus H contains C_n with S_m attached at u as a general subgraph, and so G contains $(C_n, V_2(C_n), K_m)$ as a subgraph, which is a contradiction. Therefore u is not on any cycle in H . Let C_n be any cycle in H , where $n \in \{3, \dots, \ell + 1\}$. Let P_{ℓ_0} be a shortest path between u and C_n , where $\ell_0 \in \{1, \dots, \ell\}$. Since u is not on any cycle, u has at least $d - 1 > m$ neighbors outside C_n and P_{ℓ_0} . Thus H contains $C_n \cup P_{\ell_0}$ with S_m attached at u as a general subgraph, and so G contains $(C_n, V_2(C_n), \ell_0 - 1, K_m)$ as a subgraph, which is a contradiction.

We now view H as a tree rooted at u . We claim that u is the only vertex with degree larger than m . Suppose on the contrary that there is another vertex u' with degree at least $m + 1$ in H . Let P_{ℓ_0} be a shortest path between u and u' , where $\ell_0 \in \{1, \dots, \ell\}$. Thus H contains P_{ℓ_0} with two vertex-disjoint stars S_m respectively attached to u and u' as a (general) subgraph, and so G contains $(K_m, V(K_m), \ell_0 - 1, K_m)$ as a subgraph, which is a contradiction. Therefore all the vertices but u in H has degree at most m . After the root u is removed from H , each connected component has diameter at most ℓ and degree at most m , and so each connected component has at most $m^{\ell+1} \leq N$ vertices. $\square$

The last ingredient for the proof of Theorem 1.3 for $\lambda < 2$ is the following lemma attributed to Dickson, who used it to prove a result about perfect numbers in number theory.

Lemma 2.9 (Lemma A of Dickson [13]). *For every $n \in \mathbb{N}^+$, the partially ordered set $(\mathbb{N}^n, \leq)$, in which $(a_1, \dots, a_n) \leq (b_1, \dots, b_n)$ if and only if $a_i \leq b_i$ for every i , does not contain infinite antichains.*

Dickson gave two proofs of Lemma 2.9, one of which uses induction, while the other uses Hilbert's basis theorem. For the reader's convenience, we provide a short combinatorial proof based on the infinite Ramsey's theorem.

Proof. Suppose that $s_1, s_2, \dots$ is an infinite sequence of distinct tuples in $\mathbb{N}^n$. For every $i < j$, color the edge $s_i s_j$ by 0 if $s_i < s_j$, otherwise by any $k \in \{1, \dots, n\}$ such that $s_{i,k} > s_{j,k}$. The infinite Ramsey's theorem provides an infinite subset $I \subseteq \mathbb{N}^+$ such that the edges $s_i s_j$, where $i, j \in I$ and $i \neq j$, all receive the same color c . Because $(\mathbb{N}, \leq)$ does not contain infinite descending chains, it must be the case

⁶Recall that the *diameter* of a graph is the maximum distance between a pair of vertices in the graph.

that $c = 0$. This implies that $\{s_i : i \in I\}$ is an infinite ascending chain, and in particular $\{s_1, s_2, \dots\}$ is not an antichain. $\square$

We are ready to establish the first main theorem for $\lambda < 2$.

Proof of Theorem 1.3 for $\lambda < 2$. Let $N \in \mathbb{N}$ and $\mathcal{F}_1$ be given by Lemma 2.8, and set

$$\begin{aligned}\mathcal{F}_0 &:= \{G \notin \mathcal{G}(\lambda) : G \text{ has at most } N \text{ vertices}\}, \\ \widetilde{\mathcal{F}}_2 &:= \{G \notin \mathcal{G}(\lambda) : \exists H \in \mathcal{T}_N \text{ s.t. } G = L(H)\},\end{aligned}$$

where $\mathcal{T}_N$ is the family of rooted trees such that every connected component obtained from removing the root has at most N vertices. Setting $\mathcal{F}_2$ to be the family of graphs that are minimal in $\widetilde{\mathcal{F}}_2$ under taking subgraphs, one can check that $\mathcal{F}_0 \cup \mathcal{F}_1 \cup \mathcal{F}_2$ is a forbidden subgraph characterization of $\mathcal{G}(\lambda)$.

It suffices to prove that $\mathcal{F}_2$ is finite. Let $T_1, \dots, T_n$ be an enumeration of rooted trees on at most N vertices. We encode $G \in \mathcal{F}_2$ by $t_G \in \mathbb{N}^n$ as follows. Let H be the rooted tree in $\mathcal{T}_N$ such that $G = L(H)$. After removing the root u from H , we view each connected component as a tree rooted at the vertex that is a child of u in H . Set $t_G := (t_1, \dots, t_n)$, where t_i is the number of occurrences of T_i as a connected component in the graph obtained by removing u from H . Because no member of $\mathcal{F}_2$ is a subgraph of any other, one can deduce that $\{t_G : G \in \mathcal{F}_2\}$ is an antichain in $(\mathbb{N}^n, \leq)$, and so $\mathcal{F}_2$ is finite by Lemma 2.9. $\square$

2.2. Proof of Theorem 1.3 for $\lambda \in [2, \lambda^*)$

We prove a stronger result from which the first main theorem for $\lambda \in [2, \lambda^*)$ and its corollary for $\lambda \in (2, \lambda^*)$ follow.

Theorem 2.10. *For every $\lambda \in [2, \lambda^*)$, the number of connected graphs in $\mathcal{G}(\lambda) \setminus \mathcal{G}(2)$ is finite.*

Proof of Theorem 1.3 for $\lambda \in [2, \lambda^)$ assuming Theorem 2.10.* It is known that $\mathcal{G}(2)$ has a finite forbidden subgraph characterization – in fact, there are 1,812 minimal forbidden subgraphs for $\mathcal{G}(2)$; see [6]. In view of Theorem 2.10, it suffices to show that if G is a minimal forbidden subgraph for $\mathcal{G}(\lambda)$, then G is a minimal forbidden subgraph for $\mathcal{G}(2)$, or G , after removing some vertex, is a connected graph in $\mathcal{G}(\lambda) \setminus \mathcal{G}(2)$.

Indeed, let G be a minimal forbidden subgraph for $\mathcal{G}(\lambda)$. Because $\lambda_1(G) < -\lambda \leq -2$, the graph G contains a minimal forbidden subgraph for $\mathcal{G}(2)$ as a subgraph, say F . Clearly both G and F are connected. We may assume that F is a proper subgraph of G because otherwise we are done already. Consider the graph H obtained after removing v from G , where v is a vertex of G that is furthest from F . Using the connectivity of F , one can then show that H is connected. Finally, $H \in \mathcal{G}(\lambda)$ because of the minimality of G , and $H \notin \mathcal{G}(2)$ because F is a subgraph of H . $\square$

Proof of Corollary 1.4 for $\lambda \in (2, \lambda^)$.* It follows immediately from Theorem 2.10 that $-\lambda$ is not a limit point of the set of smallest eigenvalues of graphs for $\lambda \in (2, \lambda^*)$ $\square$

The proof of Theorem 2.10 centers around the notion of *generalized line graphs*. Although we do not need their definition, we state it nevertheless for concreteness.

Definition 2.11 (Cocktail party graphs and generalized line graphs). The *cocktail party graph* $\overline{aK_2}$ is obtained from the complete graph on $2a$ vertices by deleting a perfect matching. Given a graph G with vertices $v_1, \dots, v_n$, and $a_1, \dots, a_n \in \mathbb{N}$, the *generalized line graph* $L(G; a_1, \dots, a_n)$ is obtained from the line graph $L(G)$ of G by adjoining n vertex-disjoint cocktail party graphs $\overline{a_1K_2}, \dots, \overline{a_nK_2}$ where every vertex of the i th cocktail party graph $\overline{a_iK_2}$ is adjacent to every vertex of $L(G)$ that contains v_i . See Figure 4 for an example.

Just like line graphs, all the generalized line graphs are in $\mathcal{G}(2)$, and they have a finite forbidden subgraph characterization.




Figure 4. A graph G and a schematic drawing of its generalized line graph $L(G; 2, 1, 0, 3)$.

Theorem 2.12 (Theorem 2.1 of Hoffman [17]). *The smallest eigenvalue of a generalized line graph is at least -2 .*

Theorem 2.13 (Cvetković, Doob, and Simić [10, 11], and Rao, Singhi, and Vijayan [26]). *The minimal forbidden subgraphs for the family $\mathcal{D}_\infty$ of generalized line graphs are listed in Figure 5.*

The key observation for the proof of Theorem 2.10 is that for every minimal forbidden subgraph F for $\mathcal{D}_\infty$, there exists an extension family of F disjoint from $\mathcal{G}(\lambda)$. We first deal with path extensions and clique extensions.

Lemma 2.14. *For every minimal forbidden subgraph F in Figure 5 for the family $\mathcal{D}_\infty$ of generalized line graphs, and every nonempty vertex subset A of F , the path extensions and the clique extensions of F satisfy*

$$\lim_{\ell \rightarrow \infty} \lambda_1(F, A, \ell) \leq -\lambda^* \quad \text{and} \quad \lim_{m \rightarrow \infty} \lambda_1(F, A, K_m) < -\lambda^*.$$

Moreover, the equality holds in the first inequality if and only if $F = G_4$ and $A \in \{\{3\}, \{4\}\}$.

We prove Lemma 2.14 under computer assistance in Appendix A. The next result takes care of path-clique extensions.

Lemma 2.15. *Suppose that A is a nonempty vertex subset of a graph F and $\lambda \geq 2$. If the path extensions of F satisfy*

$$\lim_{\ell \rightarrow \infty} \lambda_1(F, A, \ell) < -\lambda,$$

then there exists $m \in \mathbb{N}^+$ such that the path-clique extensions of F satisfy

$$\lambda_1(F, A, \ell, K_m) < -\lambda \text{ for every } \ell \in \mathbb{N}.$$

Proof. Pick $\ell_0 \in \mathbb{N}$ such that $\lambda_1(F, A, \ell) < -\lambda$ for every $\ell \geq \ell_0$. Clearly $\lambda_1(F, A, \ell, K_m) < -\lambda$ for every $\ell \geq \ell_0$ and every $m \in \mathbb{N}^+$. It suffices to show that for every $\ell \in \{0, \dots, \ell_0 - 1\}$ there exists $m \in \mathbb{N}^+$ such that $\lambda_1(F, A, \ell, K_m) < -\lambda$.

Let $v_0 \dots v_{\ell_0}$ be the path added to F to obtain (F, A, ℓ_0) , where the vertex v_0 is connected to every vertex in A . Set $\lambda_0 := -\lambda_1(F, A, \ell_0)$, and let $\mathbf{x}: V(F) \cup \{v_0, \dots, v_{\ell_0}\} \rightarrow \mathbb{R}$ be an eigenvector of (F, A, ℓ_0) associated with $-\lambda_0$.

Now fix $\ell \in \mathbb{N}$ with $\ell < \ell_0$, and let $m \in \mathbb{N}^+$ be determined later. Set $V_\ell := V(F) \cup \{v_0, \dots, v_\ell\}$, and identify the vertex set of (F, A, ℓ, K_m) with $V_\ell \cup V(K_m)$. We abuse notation and write x_i in place of x_{v_i} for $i \in \{\ell, \dots, \ell_0\}$. Define $\tilde{\mathbf{x}}: V_\ell \cup V(K_m) \rightarrow \mathbb{R}$ by

$$\tilde{x}_v = \begin{cases} x_v & \text{if } v \in V_\ell; \\ x_{\ell+1}/m & \text{if } v \in V(K_m). \end{cases}$$

We claim that $\sum_{v \in V_\ell} x_v^2 > 0$. Indeed, assume for the sake of contradiction that $x_v = 0$ for $v \in V_\ell$. Using $-\lambda_0 x_i = \sum_{u \sim v_i} x_u$ for $i \in \{\ell, \dots, \ell_0 - 1\}$, where the sum is taken over all vertices u that are




Figure 5. Minimal forbidden subgraphs for $\mathcal{D}_\infty$.

adjacent to the vertex v_i in (F, A, ℓ_0) , one can prove inductively that $x_i = 0$ for $i \in \{\ell + 1, \dots, \ell_0\}$, which contradicts the assumption that $\mathbf{x}$ is a nonzero vector.

Because $\sum_{v \in V_\ell} x_v^2 > 0$, clearly $\tilde{\mathbf{x}}$ is a nonzero vector. We compute

$$\tilde{\mathbf{x}}^\top \tilde{\mathbf{x}} = \sum_{v \in V_\ell} x_v^2 + m(x_{\ell+1}/m)^2.$$

Moreover we can compute $\tilde{\mathbf{x}}^\top A_{(F, A, \ell, K_m)} \tilde{\mathbf{x}}$ as follows

$$\tilde{\mathbf{x}}^\top A_{(F, A, \ell, K_m)} \tilde{\mathbf{x}} = \sum_{u, v \in V_\ell: u \sim v} x_u x_v + 2x_\ell x_{\ell+1} + m(m-1)(x_{\ell+1}/m)^2.$$

Since $\mathbf{x}$ is an eigenvector of (F, A, ℓ_0) associated with $-\lambda_0$, we obtain that

$$\sum_{u,v \in V_\ell: u \sim v} x_u x_v + x_\ell x_{\ell+1} = \sum_{v \in V_\ell} x_v \sum_{u \sim v} x_u = -\lambda_0 \sum_{v \in V_\ell} x_v^2.$$

Thus $\tilde{\mathbf{x}}^\top A_{(F,A,\ell,K_m)} \tilde{\mathbf{x}}$ can be simplified to

$$\tilde{\mathbf{x}}^\top A_{(F,A,\ell,K_m)} \tilde{\mathbf{x}} = -\lambda_0 \sum_{v \in V_\ell} x_v^2 + x_\ell x_{\ell+1} + m(m-1)(x_{\ell+1}/m)^2.$$

The Rayleigh principle says that $\lambda_1(F, A, \ell, K_m)$ is at most

$$\frac{-\lambda_0 \sum_{v \in V_\ell} x_v^2 + x_\ell x_{\ell+1} + m(m-1)(x_{\ell+1}/m)^2}{\sum_{v \in V_\ell} x_v^2 + m(x_{\ell+1}/m)^2},$$

which, as $m \rightarrow \infty$, approaches

$$-\lambda_0 + \frac{(x_\ell + x_{\ell+1})x_{\ell+1}}{\sum_{v \in V_\ell} x_v^2}.$$

Here we used the above claim that the denominator in the limit is positive.

Recall that $\lambda_0 = -\lambda_1(F, A, \ell_0) > \lambda \geq 2$. It suffices to show that $(x_\ell + x_{\ell+1})x_{\ell+1} \leq 0$. In fact, we prove inductively that $(x_i + x_{i+1})x_{i+1} \leq 0$ for $i \in \{\ell, \dots, \ell_0 - 1\}$. The base case where $i = \ell_0 - 1$ follows immediately from $-\lambda_0 x_{\ell_0} = x_{\ell_0-1}$ and $\lambda_0 > 2$. For the inductive step, using $-\lambda_0 x_{i+1} = x_i + x_{i+2}$ and $\lambda_0 > 2$, we obtain

$$\begin{aligned} (x_i + x_{i+1})x_{i+1} &= (-\lambda_0 x_{i+1} - x_{i+2} + x_{i+1})x_{i+1} = -(\lambda_0 - 2)x_{i+1}^2 - (x_{i+1} + x_{i+2})x_{i+1} \\ &\leq -(x_{i+1} + x_{i+2})x_{i+1} = -(x_{i+1} + x_{i+2})^2 + (x_{i+1} + x_{i+2})x_{i+2} \leq (x_{i+1} + x_{i+2})x_{i+2}, \end{aligned}$$

which is nonpositive by the inductive hypothesis. $\square$

We are ready to prove Theorem 2.10.

Proof of Theorem 2.10. Suppose that $\lambda \in [2, \lambda^*)$. Let $\mathcal{F}$ denote the set of minimal forbidden subgraphs for the family $\mathcal{D}_\infty$. Combining Lemmas 2.14 and 2.15, we choose $\ell, m \in \mathbb{N}^+$ such that for every $F \in \mathcal{F}$, the extension family $\mathcal{X}(F, \ell, m)$ is disjoint from $\mathcal{G}(\lambda)$. From Lemma 2.3(s), we know that $S_5 \notin \mathcal{G}(\lambda)$. In particular, no graph in $\mathcal{G}(\lambda)$ contains any member in the following family

$$\{S_5\} \cup \bigcup_{F \in \mathcal{F}} \mathcal{X}(F, \ell, m) \quad (2)$$

as a subgraph. Using Lemma 2.6, we obtain $N \in \mathbb{N}$ such that for every connected graph G with more than N vertices, if no member in (2) is a subgraph of G , then neither is any $F \in \mathcal{F}$, and so G is a generalized line graph and is in $\mathcal{G}(2)$ by Theorems 2.12 and 2.13. This implies that every connected graph in $\mathcal{G}(\lambda) \setminus \mathcal{G}(2)$ has at most N vertices. $\square$

We end this subsection with a remark on an alternative proof of Theorem 2.10. Instead of working with the family $\mathcal{D}_\infty$, one should be able to establish Lemma 2.14 for every minimal forbidden subgraph for $\mathcal{G}(2)$, and then prove Theorem 2.10 similarly. This alternative approach is more direct as it does not rely on generalized line graphs. However the drawback is obvious – there are 1,812 minimal forbidden subgraphs for $\mathcal{G}(2)$. To the best of our knowledge, there is no easily accessible database for these 1,812 graphs; for example, a complete list of these graphs was printed to microfiche in [6], and a list of only 92 of these graphs up to 8 vertices was available in [12, Fig. 2.4 and Table A1.2]. One certainly can implement the reasonably fast algorithm in [6] to enumerate the minimal forbidden subgraphs for $\mathcal{G}(2)$.

However as we strive to keep the computer-assisted part of the proof to a minimum, we work with the 31 minimal forbidden subgraphs for $\mathcal{D}_\infty$.

In fact, the family $\mathcal{G}(2)$ and its subfamily $\mathcal{D}_\infty$ are interchangeable when both families are restricted to sufficiently large connected graphs because of a characterization of $\mathcal{G}(2)$ due to Cameron, Goethals, Seidel, and Shult. To state their characterization, we introduce the following definitions.

Definition 2.16 (Representation of graphs and root systems D_n and E_8). Given a graph G and $V \subseteq \mathbb{R}^n$, we say that G is *represented* by V if the Gram matrix of V is equal to $A_G + 2I$, where A_G is the adjacency matrix of G . The *root systems* D_n and E_8 are defined by the standard basis $e_1, \dots, e_n$ of $\mathbb{R}^n$ as follows

$$D_n := \{\varepsilon_1 e_i + \varepsilon_2 e_j : \varepsilon_1, \varepsilon_2 = \pm 1, 1 \leq i < j \leq n\}, \quad E_8 := D_8 \cup \left\{ \frac{1}{2} \sum_{i=1}^8 \varepsilon_i e_i : \varepsilon_i = \pm 1, \prod_{i=1}^8 \varepsilon_i = 1 \right\}.$$

Theorem 2.17 (Theorems 4.2, 4.3 and 4.10 of Cameron et al. [7]). *For every connected graph G , the smallest eigenvalue of G is at least -2 if and only if G is represented by a subset of D_n or E_8 . Moreover*

- (a) *a graph (not necessarily connected) is represented by a subset of D_n if and only if it is a generalized line graph, and*
- (b) *a graph represented by a subset of E_8 has at most 36 vertices, and its maximum degree is at most 28.*

In particular, Theorem 2.17 implies that every connected graph in $\mathcal{G}(2)$ with more than 36 vertices is a generalized line graph.

2.3. Proof of Theorem 1.3 for $\lambda \geq \lambda^*$

Suppose that $\{F_1, \dots, F_n\}$ is a finite forbidden subgraph characterization of $\mathcal{G}(\lambda)$. Because every graph that is not in $\mathcal{G}(\lambda)$ contains F_i as a subgraph for some $i \in \{1, \dots, n\}$, no graph has its smallest eigenvalue in the open interval $(\max\{\lambda_1(F_i) : i \in \{1, \dots, n\}\}, -\lambda)$. Recall that Λ_1 consists of $\lambda \in \mathbb{R}$ such that $-\lambda$ is the smallest eigenvalue of some graph. The contrapositive of the above observation says the following.

Proposition 2.18. *Let $\lim_+ \Lambda_1 := \{\lambda \in \mathbb{R} : (\lambda, \lambda + \varepsilon) \cap \Lambda_1 \neq \emptyset \text{ for every } \varepsilon > 0\}$ be the set of right-sided limit points of Λ_1 . The family $\mathcal{G}(\lambda)$ does not have a finite forbidden subgraph characterization for any $\lambda \in \lim_+ \Lambda_1$.*

In fact, we prove that Λ_1 is dense in (λ^*, ∞) , from which the first main theorem and its corollary for $\lambda \geq \lambda^*$ follow.

Theorem 2.19. *For every $\lambda > \lambda^*$, there exist graphs $G_1, G_2, \dots$ such that $\lim_{n \rightarrow \infty} \lambda_1(G_n) = -\lambda$.*

Proof of Theorem 1.3 for $\lambda \geq \lambda^$.* Theorem 2.19 implies that $\lim_+ \Lambda_1 \supseteq [\lambda^*, \infty)$, which implies through Proposition 2.18 that $\mathcal{G}(\lambda)$ has no finite forbidden subgraph characterization for any $\lambda \geq \lambda^*$. $\square$

Proof of Corollary 1.4 for $\lambda \geq \lambda^$.* It follows immediately from Theorem 2.19 that $-\lambda$ is a limit point of the set of smallest eigenvalues of graphs for $\lambda \geq \lambda^*$. $\square$

A large chunk of Theorem 2.19 is essentially established by Shearer [28], who proved that the set of spectral radii of all graphs is dense in (λ', ∞) , where $\lambda' = \sqrt{2} + \sqrt{5}$. As was pointed out in [14], Shearer actually proved that the set of spectral radii of all caterpillar trees⁷ is dense in (λ', ∞) already. Since a caterpillar tree is bipartite, we rephrase Shearer's result in terms of smallest eigenvalues.

Theorem 2.20 (Shearer [28]; cf. Theorem 3 of Doob [14]). *For every $\lambda \geq \lambda'$, there exist caterpillar trees $G_1, G_2, \dots$ such that $\lim_{n \rightarrow \infty} \lambda_1(G_n) = -\lambda$.*

To fill the gap between $\lambda^* \approx 2.01980$ and $\lambda' \approx 2.05817$, we use the following graphs.

⁷A caterpillar tree is a tree in which all the vertices are within distance 1 of a central path.




Figure 6. A schematic drawing of the rowing graph $R(2, 0, 2, 1, 0, 8, 3, 1)$.

Definition 2.21 (Rowing graphs). Given a sequence $(a_1, \dots, a_n)$ of natural numbers, a *rowing graph* $R(a_1, \dots, a_n)$ is obtained from the path $v_{-2}v_{-1}v_0v_1 \dots v_n$ (called the *central path*) by attaching a vertex (called the *coxswain*) to v_0 , and attaching a clique of order a_i to both v_{i-1} and v_i for every $i \in \{1, \dots, n\}$. See Figure 6 for an example of a rowing graph.

We consider rowing graphs for the following two heuristics. On the one hand, a rowing graph is almost a line graph, whose smallest eigenvalue is at least -2 – when the coxswain is removed from a rowing graph, it becomes the line graph of a caterpillar tree. On the other hand, the rowing graph $R(a_1, \dots, a_n)$ contains $E_{2,n}$ (see Figure 2), whose smallest eigenvalue decreases to $-\lambda^*$ as $n \rightarrow \infty$.

We adopt the shorthand

$$(a_1, \dots, a_{k-1}, 0^{(\ell)}, a_k, \dots, a_n) := (a_1, \dots, a_{k-1}, \underbrace{0, \dots, 0}_{\ell}, a_k, \dots, a_n).$$

Lemma 2.22. *Rowing graphs satisfy the following properties.*

- (a) *The smallest eigenvalue of a rowing graph is at least $-1 - \sqrt{2}$.*
- (b) *There exists $a_1 \in \mathbb{N}$ such that $\lambda_1(R(a_1)) < -\lambda' \approx -2.05817$.*
- (c) *For every $\varepsilon > 0$ there exists $\ell \in \mathbb{N}^+$ such that*

$$\lambda_1(R(a_1, \dots, a_{n_1}, 0^{(\ell)}, b_1, \dots, b_{n_2})) > \lambda_1(R(a_1, \dots, a_{n_1}, 0^{(\ell)})) - \varepsilon$$

for every $n_1, n_2 \in \mathbb{N}$, $(a_1, \dots, a_{n_1}) \in \mathbb{N}^{n_1}$ and $(b_1, \dots, b_{n_2}) \in \mathbb{N}^{n_2}$.

- (d) *For every $n, \ell \in \mathbb{N}^+$ and $(a_1, \dots, a_n) \in \mathbb{N}^n$ with $a_n > 0$, if $\lambda_1(R(a_1, \dots, a_n, 0^{(\ell)})) \leq -2$, then for every $\varepsilon > 0$ there exists $a_{n+1} \in \mathbb{N}^+$ such that*

$$\lambda_1(R(a_1, \dots, a_{n-1}, a_n - 1, a_{n+1}, 0^{(\ell-1)})) < \lambda_1(R(a_1, \dots, a_n, 0^{(\ell)})) + \varepsilon.$$

- (e) *For every $\varepsilon > 0$, there exists $m \in \mathbb{N}^+$ such that for every $n \geq m$ and every $(a_1, \dots, a_n) \in \mathbb{N}^n$ there exists $k \in \{1, \dots, m\}$ such that*

$$\lambda_1(R(a_1, \dots, a_{k-1}, 0, a_k, \dots, a_n)) < \lambda_1(R(a_1, \dots, a_n)) + \varepsilon.$$

In the proof of Lemma 2.22, we work with vectors $\mathbf{x}$ on the vertex set of a rowing graph, whose coxswain and central path are denoted by v_c and $v_{-2}v_{-1} \dots$, and we abuse notation and write x_c and x_i in place of x_{v_c} and x_{v_i} respectively.

Proof of (a). Set $\mathbf{a} = (a_1, \dots, a_n)$ and $\lambda = \lambda_1(R(\mathbf{a}))$. Let $v_{-2} \dots v_n$ denote the central path of $R(\mathbf{a})$, and let v_c denote the coxswain attached to v_0 . Let $\mathbf{x} : V(R(\mathbf{a})) \rightarrow \mathbb{R}$ be a unit eigenvector of $R(\mathbf{a})$ associated with λ . Clearly we have $\lambda x_c = x_0$, which implies that

$$x_c^2 = \frac{x_c^2 + x_0^2}{1 + \lambda^2} \leq \frac{1}{1 + \lambda^2}. \quad (3)$$

Let L be the rowing graph $R(\mathbf{a})$ with v_c removed, and let $\mathbf{x}_L$ be $\mathbf{x}$ restricted to $V(L)$. Notice that L is a line graph of a caterpillar tree, and so $\lambda_1(L) \geq -2$ (see Footnote 1 on Page 2). Finally, we bound the smallest eigenvalue of $R(\mathbf{a})$ as follows:

$$\begin{aligned}\lambda &= \mathbf{x}^\top A_{R(\mathbf{a})} \mathbf{x} = \mathbf{x}_L^\top A_L \mathbf{x}_L + 2x_c x_0 \geq -2\mathbf{x}_L^\top \mathbf{x}_L + 2x_c x_0 \\ &= -2(1 - x_c^2) + 2\lambda x_c^2 = -2 + (2 + 2\lambda)x_c^2 \stackrel{(3)}{\geq} -2 + \frac{2 + 2\lambda}{1 + \lambda^2},\end{aligned}$$

which implies that $\lambda \geq -1 - \sqrt{2}$. In the above inequality, we assumed that $\lambda \leq -1$ which follows from the fact that $R(\mathbf{a})$ has an edge. $\square$

Proof of (b). Let $\mathbf{x}$ be the vector that assigns 1 to v_{-2} , -2 to v_{-1} , 4 to v_0 , 0 to v_1 , -2 to the coxswain, and $-4/a_1$ to every vertex in the clique of size a_1 . The Rayleigh principle says that $\lambda_1(R(a_1))$ is at most

$$\frac{\mathbf{x}^\top A_{R(a_1)} \mathbf{x}}{\mathbf{x}^\top \mathbf{x}} = \frac{-2(1 \cdot 2 + 2 \cdot 4 + 4 \cdot 2 + a_1 \cdot 4 \cdot (4/a_1)) + a_1(a_1 - 1) \cdot (-4/a_1)^2}{1^2 + (-2)^2 + 4^2 + (-2)^2 + a_1 \cdot (-4/a_1)^2},$$

which approaches $-52/25 = -2.08$ as $a_1 \rightarrow \infty$. $\square$

Proof of (c). Take $\ell = \lceil 2/\varepsilon \rceil + 5$. Let $v_{-2} \dots v_{n_1+\ell+n_2}$ denote the central path of the rowing graph $R(\mathbf{a}, 0^{(\ell)}, \mathbf{b})$, where $\mathbf{a} = (a_1, \dots, a_{n_1})$ and $\mathbf{b} = (b_1, \dots, b_{n_2})$, and let $\mathbf{x}: V(R(\mathbf{a}, 0^{(\ell)}, \mathbf{b})) \rightarrow \mathbb{R}$ be a unit eigenvector of $R(\mathbf{a}, 0^{(\ell)}, \mathbf{b})$ associated with the smallest eigenvalue. Choose $k \in \{0, \dots, \ell - 1\}$ such that $x_{n_1+k}x_{n_1+k+1}$ reaches the minimum in absolute value. In particular, using the inequality $|x_{n_1+i}x_{n_1+i+1}| \leq (x_{n_1+i}^2 + x_{n_1+i+1}^2)/2$, we obtain

$$|x_{n_1+k}x_{n_1+k+1}| \leq \frac{1}{\ell} \sum_{i=0}^{\ell-1} |x_{n_1+i}x_{n_1+i+1}| \leq \frac{1}{\ell} \sum_{i=0}^{\ell} x_{n_1+i}^2 \leq \frac{1}{\ell} < \frac{\varepsilon}{2}. \quad (4)$$

Notice that removing the edge $v_{n_1+k}v_{n_1+k+1}$ disconnects $R(\mathbf{a}, 0^{(\ell)}, \mathbf{b})$ into two subgraphs, one of which is $R(\mathbf{a}, 0^{(k)})$, while the other is a line graph, denoted L , of a caterpillar tree. Clearly

$$\lambda_1(R(\mathbf{a}, 0^{(k)})) \geq \lambda_1(R(\mathbf{a}, 0^{(\ell)})). \quad (5)$$

As $\ell \geq 6$, the graph $\widetilde{E}_8$ in Figure 1 is a proper subgraph of $R(\mathbf{a}, 0^{(\ell)})$, and so $\lambda_1(R(\mathbf{a}, 0^{(\ell)})) < -2$. Together with $\lambda_1(L) \geq -2$ (see Footnote 1 on Page 2), we obtain

$$\lambda_1(L) > \lambda_1(R(\mathbf{a}, 0^{(\ell)})). \quad (6)$$

Let $\mathbf{x}_R$ and $\mathbf{x}_L$ be the unit eigenvector $\mathbf{x}$ restricted to $V(R(\mathbf{a}, 0^{(k)}))$ and $V(L)$. Finally, we bound the smallest eigenvalue of $R(\mathbf{a}, 0^{(\ell)}, \mathbf{b})$ as follows:

$$\begin{aligned}\lambda_1(R(\mathbf{a}, 0^{(\ell)}, \mathbf{b})) &= \mathbf{x}^\top A_{R(\mathbf{a}, 0^{(\ell)}, \mathbf{b})} \mathbf{x} \\ &= \mathbf{x}_R^\top A_{R(\mathbf{a}, 0^{(k)})} \mathbf{x}_R + 2x_{n_1+k}x_{n_1+k+1} + \mathbf{x}_L^\top A_L \mathbf{x}_L \\ &\geq \lambda_1(R(\mathbf{a}, 0^{(k)}))\mathbf{x}_R^\top \mathbf{x}_R + 2x_{n_1+k}x_{n_1+k+1} + \lambda_1(L)\mathbf{x}_L^\top \mathbf{x}_L \\ &\stackrel{(4,5,6)}{>} \lambda_1(R(\mathbf{a}, 0^{(\ell)}))(\mathbf{x}_R^\top \mathbf{x}_R + \mathbf{x}_L^\top \mathbf{x}_L) - \varepsilon \\ &= \lambda_1(R(\mathbf{a}, 0^{(\ell)})) - \varepsilon.\end{aligned} \quad \square$$

Proof of (d). Set $\mathbf{a} = (a_1, \dots, a_n, 0^{(\ell)})$ and $\lambda = \lambda_1(R(\mathbf{a}))$. Let $\mathbf{x}: V(R(\mathbf{a})) \rightarrow \mathbb{R}$ be an eigenvector of $R(\mathbf{a})$ associated with λ . Let $v_{-2} \dots v_{n+\ell}$ denote the central path of the rowing graph $R(\mathbf{a})$. In $R(\mathbf{a})$, let

K denote the clique of size a_n attached to both v_{n-1} and v_n , and pick an arbitrary vertex u from K . We clearly have

$$\begin{aligned}\lambda x_u &= x_{n-1} + \sigma + x_n, \\ \lambda x_n &= x_{n-1} + \sigma + x_u + x_{n+1},\end{aligned}\tag{7}$$

where $\sigma = \sum_{u' \in V(K) \setminus \{u\}} x_{u'}$. The above identities imply that $\lambda(x_u - x_n) = x_n - x_u - x_{n+1}$, and so

$$x_u = x_n - \frac{x_{n+1}}{1 + \lambda}.\tag{8}$$

Let $R(\tilde{a})$ be the rowing graph obtained from $R(a)$ by removing u and attaching a clique, denoted $\tilde{K}$, to both v_n and v_{n+1} , where $\tilde{a} = (a_1, \dots, a_n - 1, a_{n+1}, 0^{(\ell-1)})$, and the order a_{n+1} of $\tilde{K}$ is chosen later. In particular, $V(R(\tilde{a})) = V(R(a)) \setminus \{u\} \cup V(\tilde{K})$.

We define a vector $\tilde{x}: V(R(\tilde{a})) \rightarrow \mathbb{R}$ as follows:

$$\tilde{x}_v = \begin{cases} x_n + x_u & \text{if } v = v_n; \\ -(x_n + x_u + x_{n+1})/a_{n+1} & \text{if } v \in V(\tilde{K}); \\ x_v & \text{otherwise.} \end{cases}$$

Because $\mathbf{x}$ is a nonzero vector, one can check that so is $\tilde{\mathbf{x}}$. We compute

$$\tilde{\mathbf{x}}^\top \tilde{\mathbf{x}} = \mathbf{x}^\top \mathbf{x} - x_n^2 - x_u^2 + (x_n + x_u)^2 + \frac{(x_n + x_u + x_{n+1})^2}{a_{n+1}}.$$

Moreover we can compare $\tilde{\mathbf{x}}^\top A_{R(\tilde{a})} \tilde{\mathbf{x}}$ and $\mathbf{x}^\top A_{R(a)} \mathbf{x}$ as follows

$$\begin{aligned}\tilde{\mathbf{x}}^\top A_{R(\tilde{a})} \tilde{\mathbf{x}} &= \mathbf{x}^\top A_{R(a)} \mathbf{x} - 2x_u(x_{n-1} + \sigma + x_n) + 2x_u(x_{n-1} + \sigma + x_{n+1}) \\ &\quad - 2(x_n + x_u + x_{n+1})^2 + a_{n+1}(a_{n+1} - 1) \left(\frac{x_n + x_u + x_{n+1}}{a_{n+1}} \right)^2,\end{aligned}$$

which simplifies via (7) to

$$\tilde{\mathbf{x}}^\top A_{R(\tilde{a})} \tilde{\mathbf{x}} = \lambda \mathbf{x}^\top \mathbf{x} - 2\lambda x_u^2 + 2x_u(\lambda x_n - x_u) - \left(1 + \frac{1}{a_{n+1}} \right) (x_n + x_u + x_{n+1})^2.$$

The Rayleigh principle says that $\lambda_1(R(\tilde{a}))$ is at most

$$\frac{\tilde{\mathbf{x}}^\top A_{R(\tilde{a})} \tilde{\mathbf{x}}}{\tilde{\mathbf{x}}^\top \tilde{\mathbf{x}}} = \frac{\lambda \mathbf{x}^\top \mathbf{x} - 2\lambda x_u^2 + 2x_u(\lambda x_n - x_u) - (1 + 1/a_{n+1})(x_n + x_u + x_{n+1})^2}{\mathbf{x}^\top \mathbf{x} - x_n^2 - x_u^2 + (x_n + x_u)^2 + (x_n + x_u + x_{n+1})^2/a_{n+1}},$$

which, as $a_{n+1} \rightarrow \infty$, approaches

$$\frac{\lambda \mathbf{x}^\top \mathbf{x} - 2\lambda x_u^2 + 2x_u(\lambda x_n - x_u) - (x_n + x_u + x_{n+1})^2}{\mathbf{x}^\top \mathbf{x} - x_n^2 - x_u^2 + (x_n + x_u)^2}.$$

Here we assumed that the denominator in the limit is nonzero. Indeed, suppose on the contrary that $\mathbf{x}^\top \mathbf{x} = x_n^2 + x_u^2 - (x_n + x_u)^2 = -2x_n x_u$. Because $-2x_n x_u \leq x_n^2 + x_u^2 \leq x_n^2 + x_u^2 + x_{n+1}^2 \leq \mathbf{x}^\top \mathbf{x}$, it must be the case that $x_n + x_u = 0$ and $x_{n+1} = 0$. In view of (8), we have $x_n = x_u = 0$ and hence $\mathbf{x} = \mathbf{0}$, which is a contradiction.

It suffices to prove that

$$\lambda \mathbf{x}^\top \mathbf{x} - 2\lambda x_u^2 + 2x_u(\lambda x_n - x_u) - (x_n + x_u + x_{n+1})^2 \leq \lambda (\mathbf{x}^\top \mathbf{x} - x_n^2 - x_u^2 + (x_n + x_u)^2),$$

which is equivalent to

$$(x_n + x_u + x_{n+1})^2 + 2(1 + \lambda)x_u^2 \geq 0.$$

Using (8), we know that the left hand side of the last inequality is equal to

$$\left(2x_n + \frac{\lambda}{1+\lambda}x_{n+1}\right)^2 + 2(\lambda+1)\left(x_n - \frac{x_{n+1}}{1+\lambda}\right)^2 = (2\lambda+6)x_n^2 - \frac{4}{1+\lambda}x_nx_{n+1} + \left(1 + \frac{1}{(1+\lambda)^2}\right)x_{n+1}^2.$$

From Lemma 2.22(a), we know that $\lambda \geq -1 - \sqrt{2}$ in addition to the assumption that $\lambda \leq -2$. One can check

$$1 + \frac{1}{(1+\lambda)^2} \geq 2\lambda+6 \geq \frac{4}{1+\lambda} \cdot \frac{1}{\lambda} > 0, \quad \text{for } \lambda \in [-1 - \sqrt{2}, -2].$$

We are left to prove $x_n^2 - \lambda x_n x_{n+1} + x_{n+1}^2 \geq 0$. In fact, we prove inductively that $x_i^2 - \lambda x_i x_{i+1} + x_{i+1}^2 \geq 0$ for $i \in \{n, \dots, n + \ell - 1\}$. The base case where $i = n + \ell - 1$ follows immediately from $\lambda x_{n+\ell} = x_{n+\ell-1}$. For the inductive step, using $\lambda x_{i+1} = x_i + x_{i+2}$, we obtain

$$x_i^2 - \lambda x_i x_{i+1} + x_{i+1}^2 = (\lambda x_{i+1} - x_{i+2})^2 - \lambda(\lambda x_{i+1} - x_{i+2})x_{i+1} + x_{i+1}^2 = x_{i+1}^2 - \lambda x_{i+1} x_{i+2} + x_{i+2}^2,$$

which is nonnegative by the inductive hypothesis. $\square$

Proof of (e). Take $m = \lceil 5/\varepsilon \rceil + 1$. Let $v_{-2} \dots v_n$ denote the central path of the rowing graph $R(\mathbf{a})$, where $\mathbf{a} = (a_1, \dots, a_n)$, and let $\mathbf{x} : V(R(\mathbf{a})) \rightarrow \mathbb{R}$ be a unit eigenvector associated with the smallest eigenvalue of $R(\mathbf{a})$. Choose $k \in \{1, \dots, m\}$ such that x_{k-1} reaches the minimum in absolute value. In particular,

$$x_{k-1}^2 \leq \frac{1}{m} \sum_{i=0}^{m-1} x_i^2 \leq \frac{1}{m} < \frac{\varepsilon}{5}. \quad (9)$$

Let $v_{-2} \dots v_{k-1} v_* v_k \dots v_n$ denote the central path of $R(\tilde{\mathbf{a}})$, where $\tilde{\mathbf{a}} = (a_1, \dots, a_{k-1}, 0, a_k, \dots, a_n)$. We naturally view the vertex set of $R(\tilde{\mathbf{a}})$ as $V(R(\mathbf{a})) \cup \{v_*\}$, and we extend the unit eigenvector $\mathbf{x} : V(R(\mathbf{a})) \rightarrow \mathbb{R}$ to $\tilde{\mathbf{x}} : V(R(\tilde{\mathbf{a}})) \rightarrow \mathbb{R}$ by setting $\tilde{x}_* = x_{k-1}$. The Rayleigh principle says that $\lambda_1(R(\tilde{\mathbf{a}}))$ is at most

$$\begin{aligned} \frac{\tilde{\mathbf{x}}^\top A_{R(\tilde{\mathbf{a}})} \tilde{\mathbf{x}}}{\tilde{\mathbf{x}}^\top \tilde{\mathbf{x}}} &= \frac{\mathbf{x}^\top A_{R(\mathbf{a})} \mathbf{x} + 2x_{k-1}^2}{\mathbf{x}^\top \mathbf{x} + x_{k-1}^2} = \frac{\lambda_1(R(\mathbf{a})) + 2x_{k-1}^2}{1 + x_{k-1}^2} \\ &= \lambda_1(R(\mathbf{a})) + \frac{(2 - \lambda_1(R(\mathbf{a})))x_{k-1}^2}{1 + x_{k-1}^2} \leq \lambda_1(R(\mathbf{a})) + (2 - \lambda_1(R(\mathbf{a})))x_{k-1}^2. \end{aligned}$$

From Lemma 2.22(a), we obtain

$$(2 - \lambda_1(R(\mathbf{a})))x_{k-1}^2 \leq (2 + (1 + \sqrt{2}))x_{k-1}^2 \stackrel{(9)}{<} \varepsilon. \quad \square$$

We now have all of the ingredients needed to establish Theorem 2.19.

Proof of Theorem 2.19. In view of Theorem 2.20, we may assume that $\lambda \in (\lambda^*, \lambda')$. Fix $\varepsilon > 0$. We assume for the sake of contradiction that no rowing graph has its smallest eigenvalue in $(-\lambda - \varepsilon, -\lambda)$ because otherwise we are done. Define

$$\begin{aligned} S &:= \{(a_1, \dots, a_n) \in \mathbb{N}^n : n \in \mathbb{N}^+ \text{ and } a_n > 0\}, \\ A &:= \{(a_1, \dots, a_n) \in S : \lambda_1(R(a_1, \dots, a_n, 0^{(\ell)})) < -\lambda \text{ for some } \ell \in \mathbb{N}\}. \end{aligned}$$

Claim. For every $m \in \mathbb{N}^+$ there exists $(a_1, \dots, a_n) \in A$ with exactly m nonzero entries such that

$$(a_1, \dots, a_{k-1}, 0, a_k, \dots, a_n) \notin A \text{ for every } k \in \{1, \dots, n\}. \quad (10)$$

Proof of Claim. We order the elements in S as follows:

$(a_1, \dots, a_{n_1}) < (b_1, \dots, b_{n_2})$ if and only if

$(a_1, \dots, a_{n_1}, 0^{(n_2)})$ strictly precedes $(b_1, \dots, b_{n_2}, 0^{(n_1)})$ lexicographically.

Alternatively, $(a_1, \dots, a_{n_1}) < (b_1, \dots, b_{n_2})$ if and only if there exists $\delta > 0$ such that $\sum_{i=1}^{n_1} a_i x^i < \sum_{i=1}^{n_2} b_i x^i$ for every $x \in (0, \delta)$. We shall repeatedly use the fact that for every $\ell \in \mathbb{N}^+$ the set

$$S_\ell := \{a \in S : \text{the length of } a \text{ is at most } \ell\}$$

does not contain an infinite descending chain, hence $(S_\ell, \leq)$ is well-founded, that is, every nonempty subset of S_ℓ has a minimal element.⁸ This fact can be established by a simple induction on ℓ .

In particular, the ordering on S implies that $(a_1, \dots, a_{k-1}, 0, a_k, \dots, a_n) < (a_1, \dots, a_n)$ when both sequences are in S and $k \in \{1, \dots, n\}$. Thus it suffices to construct, for every $m \in \mathbb{N}^+$, a minimal element $a^{(m)}$ of A_m , where

$$A_m := \{(a_1, \dots, a_n) \in A : (a_1, \dots, a_n) \text{ has } m \text{ nonzero entries}\}.$$

Our inductive construction additionally requires that $a^{(1)}, a^{(2)}, a^{(3)}, \dots$ form a descending chain.

Apply Lemma 2.22(c) to obtain $\ell \in \mathbb{N}^+$ such that

$$\begin{aligned} \lambda_1(R(a_1, \dots, a_{n_1}, 0^{(\ell)}, b_1, \dots, b_{n_2})) &> \lambda_1(R(a_1, \dots, a_{n_1}, 0^{(\ell)})) - \varepsilon \\ &\text{for every } n_1, n_2 \in \mathbb{N}, (a_1, \dots, a_{n_1}) \in \mathbb{N}^{n_1} \text{ and } (b_1, \dots, b_{n_2}) \in \mathbb{N}^{n_2}. \end{aligned} \quad (11)$$

In particular, when $n_1 = 0$, we have

$$\lambda_1(R(0^{(\ell)}, b_1, \dots, b_{n_2})) > \lambda_1(R(0^{(\ell)})) - \varepsilon \text{ for every } (b_1, \dots, b_{n_2}) \in \mathbb{N}^{n_2}.$$

For the base case where $m = 1$, note that $R(0^{(\ell)})$ is just $E_{2,\ell}$ defined in Proposition 2.1, which satisfies that

$$\lambda_1(R(0^{(\ell)})) = \lambda_1(E_{2,\ell}) > -\lambda^* > -\lambda.$$

Since no rowing graph has its smallest eigenvalue in $(-\lambda - \varepsilon, -\lambda)$, the above two inequalities imply

$$\lambda_1(R(0^{(\ell)}, b_1, \dots, b_{n_2})) \geq -\lambda \text{ for every } (b_1, \dots, b_{n_2}) \in \mathbb{N}^{n_2},$$

which further implies that the length of every sequence in A_1 is at most ℓ , that is, $A_1 \subseteq S_\ell$. Moreover, Lemma 2.22(b) implies that A_1 is nonempty. Therefore there exists a minimal element of A_1 .

For the inductive step, suppose that $m \in \mathbb{N}^+$, and that $a^{(1)} > \dots > a^{(m)}$ are minimal elements of $A_1, \dots, A_m$ respectively. Say $a^{(m)} = (b_1, \dots, b_n)$. Lemma 2.22(d) shows that there exists $b_{n+1} \in \mathbb{N}^+$ such that $b^{(m+1)} := (b_1, \dots, b_{n-1}, b_n - 1, b_{n+1}) \in A$. Since $b^{(m+1)} < a^{(m)}$ and $a^{(m)}$ is minimal in A_m , it must be the case that $b_n > 1$ or equivalently $b^{(m+1)} \in A_{m+1}$. It suffices to show that

$$\{b \in A_{m+1} : b \leq b^{(m+1)}\} \subseteq S_{(m+1)\ell}.$$

⁸Assuming the axiom of dependent choice, a weak form of the axiom of choice, $(S_\ell, \leq)$ is well-founded if it contains no countable infinite descending chains.

Assume for the sake of contradiction that there exists $\mathbf{b} \in A_{m+1}$ such that $\mathbf{b} \leq \mathbf{b}^{(m+1)}$ and the length of $\mathbf{b}$ is more than $(m+1)\ell$. By the pigeonhole principle, there exists an initial segment, denoted $\mathbf{b}'$, of $\mathbf{b}$ such that $(\mathbf{b}', 0^{(\ell)})$ is an initial segment of $\mathbf{b}$. We may require in addition that either $\mathbf{b}'$ is the empty sequence or the last entry of $\mathbf{b}'$ is positive. If $\mathbf{b}'$ is the empty sequence, then we can argue similarly to the base case that

$$\lambda_1(R(\mathbf{b}, c_1, \dots, c_{n'})) \geq -\lambda \text{ for every } (c_1, \dots, c_{n'}) \in \mathbb{N}^{n'},$$

which contradicts $\mathbf{b} \in A_{m+1}$.

We may assume that the last entry of $\mathbf{b}'$ is positive. Because $(\mathbf{b}', 0^{(\ell)})$ is an initial segment of $\mathbf{b}$, and the last entry of $\mathbf{b}$ is positive, the number of nonzero entries in $\mathbf{b}'$, denoted $i \in \mathbb{N}^+$, is at most m . Since $\mathbf{b}' < \mathbf{b} \leq \mathbf{b}^{(m+1)} < \mathbf{a}^{(m)} \leq \mathbf{a}^{(i)}$, and $\mathbf{a}^{(i)}$ is minimal in A_i , we know that $\mathbf{b}' \notin A$, and in particular

$$\lambda_1(R(\mathbf{b}', 0^{(\ell)})) \geq -\lambda.$$

From (11), we know that

$$\lambda_1(R(\mathbf{b}, c_1, \dots, c_{n'})) > \lambda_1(R(\mathbf{b}', 0^{(\ell)})) - \varepsilon \text{ for every } (c_1, \dots, c_{n'}) \in \mathbb{N}^{n'}.$$

Since no rowing graph has its smallest eigenvalue in $(-\lambda - \varepsilon, -\lambda)$, the above two inequalities imply

$$\lambda_1(R(\mathbf{b}, c_1, \dots, c_{n'})) \geq -\lambda \text{ for every } (c_1, \dots, c_{n'}) \in \mathbb{N}^{n'},$$

which contradicts $\mathbf{b} \in A_{m+1}$. □

Finally, let m be given by Lemma 2.22(e). The claim provides $(a_1, \dots, a_n) \in A$ with m nonzero entries such that (10) holds. Let $\ell \in \mathbb{N}$ be such that

$$\lambda_1(R(a_1, \dots, a_n, 0^{(\ell)})) < -\lambda.$$

Since the length of $(a_1, \dots, a_n, 0^{(\ell)})$ is at least m , Lemma 2.22(e) says that there exists $k \in \{1, \dots, m\}$ such that

$$\lambda_1(R(a_1, \dots, a_{k-1}, 0, a_k, \dots, a_n, 0^{(\ell)})) < \lambda_1(R(a_1, \dots, a_n, 0^{(\ell)})) + \varepsilon,$$

However (10) asserts that $(a_1, \dots, a_{k-1}, 0, a_k, \dots, a_n) \notin A$, which implies that

$$\lambda_1(R(a_1, \dots, a_{k-1}, 0, a_k, \dots, a_n, 0^{(\ell)})) \geq -\lambda.$$

Combining the last three inequalities, we obtain that

$$-\lambda - \varepsilon < \lambda_1(R(a_1, \dots, a_n, 0^{(\ell)})) < -\lambda. \quad \square$$

3. Forbidden subgraphs for $\mathcal{G}^\pm(\lambda)$

A useful tool in spectral graph theory for signed graphs is switching – two signed graphs are *switching equivalent* if one graph can be obtained from the other by reversing all the edges in a cut-set. An important feature of switching equivalence is that the switching equivalent signed graphs all have the same spectrum.

Hereinafter we adopt the following convention for an unsigned graph G . With a slight abuse of notation, we denote by G the all-positive signed graph with underlying graph G . We also denote by $-G$ the all-negative signed graph with the same underlying graph.

To prove the second main theorem, we need to extend the concepts and results from the previous section. We recommend that the reader go through the rest of the section alongside Section 2.

3.1. Proof of Theorem 1.5 for $\lambda < 2$

We first generalize path extensions, path-clique extensions, clique extensions, and Lemma 2.3(c1, c2).

Definition 3.1. Given a nonempty vertex subset A of a signed graph $F^\pm$, a *signed vertex subset* $A^\pm$ of $F^\pm$ is the vertex subset A together with an assignment of signs (positive or negative). Given, in addition, $\ell \in \mathbb{N}$ and $m \in \mathbb{N}^+$,

- (a) the *path extension* $(F^\pm, A^\pm, \ell)$ is obtained from $F^\pm$ by adding an all-positive path $v_0 \dots v_\ell$ of length ℓ and connecting v_0 to every vertex v in A by an edge signed according to the sign of v in $A^\pm$;
- (b) the *path-clique extension* $(F^\pm, A^\pm, \ell, K_m)$ is further obtained from $(F^\pm, A^\pm, \ell)$ by adding an all-positive clique of size m and connecting every vertex in the clique to v_ℓ by a positive edge;
- (c) the *clique extension* $(F^\pm, A^\pm, K_m)$ is obtained from $F^\pm$ by adding a clique of order m and connecting every vertex in the clique to every vertex v in A by an edge signed according to the sign of v in $A^\pm$.

Lemma 3.2. For every signed cycle $C_n^\pm$ of length n and every signed set $A^\pm$ of two adjacent vertices of $C_n^\pm$, the path-clique extensions and the clique extensions of $C_n^\pm$ satisfy:

- (c1) $\lim_{m \rightarrow \infty} \lambda_1(C_n^\pm, A^\pm, \ell, K_m) \leq -2$ for fixed $\ell \in \mathbb{N}$;
- (c2) $\lim_{m \rightarrow \infty} \lambda_1(C_n^\pm, A^\pm, K_m) \leq -2$.

Proof. Let $v_0, v_1, \dots, v_{n-1}$ be the vertices of $C_n^\pm$, and let $\sigma: E(C_n^\pm) \rightarrow \{\pm 1\}$ be the signing of $C_n^\pm$. Suppose that $A^\pm$ is a signed set of $\{v_0, v_{n-1}\}$. By switching a suitable subset of $\{v_0, v_{n-1}\}$, we may assume that $A^\pm = \{v_0^+, v_{n-1}^+\}$. If $C_n^\pm$ has an even number of positive edges, then $C_n^\pm$ is already switching equivalent to the all-negative cycle of length n , whose smallest eigenvalue is -2 . If the edge $v_0 v_{n-1}$ is negative, then both $(C_n^\pm, A^\pm, \ell, K_m)$ and $(C_n^\pm, A^\pm, K_m)$ contain a subgraph that is switching equivalent to the all-negative triangle $-K_3$, and so their smallest eigenvalues are at most $\lambda_1(-K_3) = -2$.

Hereafter we assume that $C_n^\pm$ has an odd number of positive edges and the edge $v_0 v_{n-1}$ is positive. Furthermore, by switching a suitable subset of $\{v_1, \dots, v_{n-2}\}$, we may assume that $v_0 v_{n-1}$ is the only positive edge of the signed cycle $C_n^\pm$.

For (c1), we define a vector $\mathbf{x}: V(C_n^\pm, A^\pm, \ell, K_m) \rightarrow \mathbb{R}$ as follows. Let $v_n v_{n+1} \dots v_{n+\ell}$ and K_m be the path and the clique added to $C_n^\pm$ to obtain the path-clique extension $(C_n^\pm, A^\pm, \ell, K_m)$, where v_n is adjacent to v_0 and v_{n-1} , and $v_{n+\ell}$ is adjacent to every vertex in K_m . Assign $x_{v_i} = -1$ for $i \in \{0, \dots, n-1\}$, $x_{v_{n+i}} = 2(-1)^i$ for $i \in \{0, \dots, \ell\}$, and $x_u = -x_{n+\ell}/m$ for every $u \in V(K_m)$. We abuse notation and write x_i in place of x_{v_i} for $i \in \{0, \dots, n+\ell\}$. The Rayleigh principle says that the smallest eigenvalue of $(C_n^\pm, A^\pm, \ell, K_m)$ is at most

$$\left(-2 \sum_{i=0}^{n-2} x_i x_{i+1} + 2x_0 x_{n-1} + 2(x_0 + x_{n-1})x_n + 2 \sum_{i=0}^{\ell-1} x_{n+i} x_{n+i+1} - 2x_{n+\ell}^2 + m(m-1)(x_{n+\ell}/m)^2 \right) / \left(\sum_{i=0}^{n-1} x_i^2 + \sum_{i=0}^{\ell} x_{n+i}^2 + m(x_{n+\ell}/m)^2 \right),$$

which is equal to $(-2n - 8(\ell+1) - 4/m)/(n + 4(\ell+1) + 4/m)$, and approaches -2 as $m \rightarrow \infty$.

For (c2), we define a vector $\mathbf{x}: V(C_n^\pm, A^\pm, K_m) \rightarrow \mathbb{R}$ as follows. Let K_m be the clique added to $C_n^\pm$ to obtain the clique extension $(C_n^\pm, A^\pm, K_m)$. Assign $x_{v_i} = -1$ for $i \in \{0, \dots, n-1\}$ and $x_u = -(x_0 + x_{n-1})/m$ for every $u \in V(K_m)$. We abuse notation and write x_i in place of x_{v_i} for $i \in \{0, \dots, n-1\}$. The Rayleigh principle says that the smallest eigenvalue of $(C_n^\pm, A^\pm, K_m)$ is at most

$$\frac{-2 \sum_{i=0}^{n-2} x_i x_{i+1} + 2x_0 x_{n-1} - 2(x_0 + x_{n-1})^2 + m(m-1)((x_0 + x_{n-1})/m)^2}{\sum_{i=0}^{n-1} x_i^2 + m((x_0 + x_{n-1})/m)^2},$$

which is equal to $(-2n - 4/m)/(n + 4/m)$, and approaches -2 as $m \rightarrow \infty$. □

We naturally generalize extension families as well as Lemma 2.5.

Definition 3.3 (Signed extension family). Given a signed graph $F^\pm$ and $\ell, m \in \mathbb{N}^+$, the *signed extension family* $\mathcal{X}^\pm(F^\pm, \ell, m)$ of $F^\pm$ consists of the path-extension $(F^\pm, A^\pm, \ell)$, the path-clique extensions $(F^\pm, A^\pm, \ell_0, K_m)$, and the clique extension $(F^\pm, A^\pm, K_m)$, where $A^\pm$ ranges over the nonempty signed vertex subsets of $F^\pm$, and ℓ_0 ranges over $\{0, \dots, \ell - 1\}$.

Note that for an unsigned graph F and $\ell, m \in \mathbb{N}^+$, the extension family $\mathcal{X}(F, \ell, m)$ (defined in Definition 2.4) is a subfamily of the signed extension family $\mathcal{X}^\pm(F, \ell, m)$.

Lemma 3.4. *For every $\lambda < 2$, there exist $\ell, m \in \mathbb{N}^+$ such that both the signed extension family $\mathcal{X}^\pm(C, \ell, m)$ of the claw graph C and the signed extension family $\mathcal{X}^\pm(D, \ell, m)$ of the diamond graph D are disjoint from $\mathcal{G}^\pm(\lambda)$.*

Proof. From Lemma 2.3(n1, n2, p) and Lemma 2.5, we obtain $\ell, m \in \mathbb{N}^+$ such that $P_\ell \notin \mathcal{G}^\pm(\lambda)$, none of the following graphs

$$(\overline{K}_2, V(\overline{K}_2), 0, K_m), \dots, (\overline{K}_2, V(\overline{K}_2), \ell - 1, K_m), \text{ and } (\overline{K}_2, V(\overline{K}_2), K_m) \quad (12)$$

is in $\mathcal{G}^\pm(\lambda)$, and $\mathcal{X}(C, \ell, m) \cup \mathcal{X}(D, \ell, m)$ is disjoint from $\mathcal{G}^\pm(\lambda)$. Since $P_\ell \notin \mathcal{G}^\pm(\lambda)$, no path extension in $\mathcal{X}^\pm(C, \ell, m) \cup \mathcal{X}^\pm(D, \ell, m)$ is in $\mathcal{G}^\pm(\lambda)$.

We are left to deal with the path-clique extensions and the clique extension

$$(F, A^\pm, 0, K_m), \dots, (F, A^\pm, \ell - 1, K_m), \text{ and } (F, A^\pm, K_m), \quad (13)$$

where F is C or D , and $A^\pm$ is the signed set of a nonempty vertex subset A of F . We break the rest of the proof into three cases.

Case 1: $A^\pm$ is all-positive or all-negative. This case follows since the signed graphs in (13), after switching the cut-set between $V(F)$ and its complement in case $A^\pm$ is all-negative, become respectively $(F, A, 0, K_m), \dots, (F, A, \ell - 1, K_m)$, and (F, A, K_m) from the extension family $\mathcal{X}(F, \ell, m)$, which is disjoint from $\mathcal{G}^\pm(\lambda)$.

Case 2: There exists an edge uv of F such that $\{u^-, v^+\} \subseteq A^\pm$. Since the all-negative triangle $-K_3$, whose smallest eigenvalue is -2 , is switching equivalent to a subgraph of $(F, A^\pm, 0)$, no signed graph in (13) is in $\mathcal{G}^\pm(\lambda)$.

Case 3: $A^\pm$ is neither all-positive nor all-negative, and no edge uv of F satisfies that $\{u^-, v^+\} \subseteq A^\pm$. Label the vertices of C and D as in Figure 3. We may assume without loss of generality that $\{1^-, 3^+\} \subseteq A^\pm$. One can check that, for $F \in \{C, D\}$, the signed graphs in (12), after switching at the vertex labeled by 1, are respectively subgraphs of those in (13), and hence no signed graph in (13) is in $\mathcal{G}^\pm(\lambda)$. $\square$

Next, generalizing Lemma 2.6, we show that forbidding a star, an all-negative complete graph, and a signed extension family of $F^\pm$ effectively forbids $F^\pm$ itself in every sufficiently large connected signed graph.

Lemma 3.5. *For every signed graph $F^\pm$ and $k_1, k_2, \ell, m \in \mathbb{N}^+$, there exists $N \in \mathbb{N}$ such that for every connected signed graph $G^\pm$ with more than N vertices, if $G^\pm$ does not contain any subgraph that is switching equivalent to any member in $\{S_{k_1}, -K_{k_2}\} \cup \mathcal{X}^\pm(F^\pm, \ell, m)$, then $G^\pm$ does not contain any subgraph that is switching equivalent to $F^\pm$ either.*

We leave the proof to the reader as one can make the proof of Lemma 2.6 into that for Lemma 3.5 by *mutatis mutandis*. To get started, one might want to take $N = vd^\ell$, where d is the Ramsey number $R(k_1, k_2, 3^{v+1}m + v)$.

The last ingredient is a sufficient condition for signed line graphs, which generalizes Theorem 2.7.

Definition 3.6 (Bidirected graph and signed line graph). A *bidirected graph* is a graph in which each vertex-edge incidence has a positive or negative sign. We decorate variables for bidirected graphs with an arrow on the top. A signed graph $G^\pm$ is the *signed line graph* of a bidirected graph $\vec{H}$ if the underlying




Figure 7. A bidirected graph and its signed line graph. In a signed graph, the positive edges are represented by solid segments and the negative edges are represented by dashed segments.

graph of $G^\pm$ is the line graph of $\vec{H}$, and moreover for every two distinct edges e_1 and e_2 of $\vec{H}$ having a vertex v in common, the sign of the edge e_1e_2 in $G^\pm$ is the product of the signs of the two incidences (v, e_1) and (v, e_2) in $\vec{H}$.

Pictorially we place a sign close to each incidence in a bidirected graph. See Figure 7 for an example of a bidirected graph and its signed line graph.

Remark. Our definition of a signed line graph is different from the existing literature – the conventional definition (e.g., Zaslavsky [34]) reverses all the edge signs of $G^\pm$ in Definition 3.6. Our choice of the definition is based on aesthetic reasons: the signed line graph of an all-positive bidirected graph is still all-positive, and the smallest eigenvalue of a signed line graph is at least -2 .

Lemma 3.7 (Proposition 2.2 of Vijayakumar [31]). *For every signed graph $G^\pm$, if $G^\pm$ does not contain any subgraph that is switching equivalent to the claw graph, the diamond graph, or the all-negative triangle, then $G^\pm$ is a signed line graph.*

Although the conclusion in [31, Proposition 2.2] is weaker than that in Lemma 3.7, Vijayakumar's proof already gives Lemma 3.7. Below we reproduce his proof in terms of signed line graphs.

Proof. Let G be the underlying graph of $G^\pm$. One can check that G contains neither the claw graph nor the diamond graph as a subgraph. Thus Theorem 2.7 implies that G is a line graph of another graph, denoted H , without isolated vertices. We may identify the vertex set of $G^\pm$ with the edge set of H .

For every $v \in V(H)$, let $E_v(H)$ be the set of edges that are incident to v in H . Define a signing $\sigma: \{(v, e): v \in V(H), e \in E_v(H)\} \rightarrow \{\pm 1\}$ such that for every $v \in V(H)$,

$$\sigma(v, e_1)\sigma(v, e_2) = A_{G^\pm}(e_1, e_2) \text{ for distinct } e_1, e_2 \in E_v(H). \quad (14)$$

Such a signing σ can be chosen as follows: for every $v \in V(H)$, define $\sigma(v, e_0) = 1$ for some arbitrary $e_0 \in E_v(H)$, and define $\sigma(v, e) = A_{G^\pm}(e_0, e)$ for any other $e \in E_v(H)$. Since $G^\pm$ does not contain any subgraph that is switching equivalent to the all-negative triangle, it can be seen that (14) holds. Assigning $\sigma(v, e)$ to every incidence (v, e) of H , we obtain a bidirected graph $\vec{H}$ whose signed line graph is $G^\pm$. $\square$

Similar to Lemma 2.8, we obtain a finite set of forbidden subgraphs for $\mathcal{G}^\pm(\lambda)$ that forces every sufficiently large connected signed graph to be the signed line graph of a bidirected tree whose complexity is uniformly bounded.

Lemma 3.8. *For every $\lambda < 2$, there exist $N \in \mathbb{N}$ and a finite family $\mathcal{F}_1^\pm$ that is disjoint from $\mathcal{G}^\pm(\lambda)$ such that for every connected signed graph $G^\pm$ with more than N vertices, if $G^\pm$ contains no member in $\mathcal{F}_1^\pm$ as a subgraph, then there exists a bidirected rooted tree $\vec{H} \in \vec{\mathcal{T}}_N$ such that $G^\pm$ is the signed line graph of $\vec{H}$, where $\vec{\mathcal{T}}_N$ is the family of bidirected rooted tree such that every connected component obtained from removing the root has at most N vertices.*

Proof. Denote $\mathcal{C}_n^\pm$ the set of signed cycles of length n . We obtain $\ell, m \in \mathbb{N}^+$ from Lemmas 2.3, 3.2 and 3.4 such that the following family

$$\begin{aligned}\tilde{\mathcal{F}}_1^\pm &:= \{S_4, P_\ell, -K_3\} \cup \mathcal{X}^\pm(C, \ell, m) \cup \mathcal{X}^\pm(D, \ell, m) \\ &\cup \{(C_n^\pm, V_2(C_n), \ell_0, K_m): n \in \{3, \dots, \ell + 1\}, C_n^\pm \in \mathcal{C}_n^\pm, \ell_0 \in \{0, \dots, \ell - 1\}\} \\ &\cup \{(C_n^\pm, V_2(C_n), K_m): n \in \{3, \dots, \ell + 1\}, C_n^\pm \in \mathcal{C}_n^\pm\} \\ &\cup \{(K_m, V(K_m), \ell_0, K_m): \ell_0 \in \{0, \dots, \ell - 1\}\}\end{aligned}$$

is disjoint from $\mathcal{G}^\pm(\lambda)$. Let $\mathcal{F}_1^\pm$ be the smallest family, that is closed under switching, containing $\tilde{\mathcal{F}}_1^\pm$. Clearly $\mathcal{F}_1^\pm$ is also disjoint from $\mathcal{G}^\pm(\lambda)$.

We omit the rest of the proof as it, *mutatis mutandis*, is very much like the one in the proof of Theorem 1.3 for $\lambda < 2$ on Page 7. We point out that one should make use of Lemmas 3.5 and 3.7 in place of Lemma 2.6 and Theorem 2.7 respectively. $\square$

We are ready to present the proof of the second main theorem for $\lambda < 2$.

Proof of Theorem 1.5 for $\lambda < 2$. Let $N \in \mathbb{N}$ and $\mathcal{F}_1^\pm$ be given by Lemma 3.8, and set

$$\begin{aligned}\mathcal{F}_0^\pm &:= \{G^\pm \notin \mathcal{G}^\pm(\lambda): G^\pm \text{ has at most } N \text{ vertices}\}, \\ \tilde{\mathcal{F}}_2^\pm &:= \{G^\pm \notin \mathcal{G}^\pm(\lambda): \exists \vec{H} \in \vec{\mathcal{T}}_N \text{ s.t. } G^\pm \text{ is the signed line graph of } \vec{H}\},\end{aligned}$$

where $\vec{\mathcal{T}}_N$ is the family of bidirected rooted tree such that every connected component obtained from removing the root has at most N vertices. Setting $\mathcal{F}_2^\pm$ to be the family of signed graphs that are minimal in $\tilde{\mathcal{F}}_2^\pm$ under taking subgraphs, one can check that $\mathcal{F}_0^\pm \cup \mathcal{F}_1^\pm \cup \mathcal{F}_2^\pm$ is a forbidden subgraph characterization of $\mathcal{G}^\pm(\lambda)$.

It suffices to prove that $\mathcal{F}_2^\pm$ is finite. Let $\vec{T}_1, \dots, \vec{T}_n$ be an enumeration of bidirected rooted trees $\vec{T}$ on at most $N + 1$ vertices. We encode $G^\pm \in \mathcal{F}_2^\pm$ by $t_{G^\pm} \in \mathbb{N}^n$ as follows. Let $\vec{H}$ be the bidirected rooted tree in $\vec{\mathcal{T}}_N$ such that $G^\pm$ is the signed line graph of $\vec{H}$. After removing the root u from $\vec{H}$, let $U_1, U_2, \dots, U_m$ be the vertex sets of the connected components. For $i \in \{1, \dots, m\}$, we view the subgraph $\vec{H}_i := \vec{H}[\{u\} \cup U_i]$ as a bidirected tree rooted at u . Set $t_{G^\pm} = (t_1, \dots, t_n)$, where t_i is the number of occurrences of $\vec{T}_i$ in $\vec{H}_1, \dots, \vec{H}_m$. Because no member of $\mathcal{F}_2^\pm$ is a subgraph of any other, one can deduce that $\{t_{G^\pm}: G^\pm \in \mathcal{F}_2^\pm\}$ is an antichain in $(\mathbb{N}^n, \leq)$, and so $\mathcal{F}_2^\pm$ is finite by Lemma 2.9. $\square$

3.2. Proof of Theorem 1.5 for $\lambda \in [2, \lambda^*)$

We prove a generalization of Theorem 2.10, from which the second main theorem for $\lambda \in [2, \lambda^*)$ follows.

Theorem 3.9. *For every $\lambda \in [2, \lambda^*)$, the number of connected signed graphs in $\mathcal{G}^\pm(\lambda) \setminus \mathcal{G}^\pm(2)$ is finite.*

Proof of Theorem 1.5 for $\lambda \in [2, \lambda^)$.* Using the fact that every minimal forbidden subgraph for $\mathcal{G}^\pm(2)$ has at most 10 vertices (see [31]), and Theorem 3.9, one can prove this case by following the argument in the proof of Theorem 1.3 for $\lambda \in [2, \lambda^*)$ on Page 9. $\square$

Recall from Theorem 2.17(a) that a graph is a generalized line graph if and only if it is represented by a subset of the root system D_n defined in Definition 2.16. For signed graphs, we work directly with those that are represented by a subset of D_n .

Definition 3.10 (Representation of signed graphs). Given a signed graph $G^\pm$ and a subset V of $\mathbb{R}^n$, we say that $G^\pm$ is *represented* by V if the Gram matrix of V is equal to $A_{G^\pm} + 2I$, where $A_{G^\pm}$ is the signed adjacency matrix of $G^\pm$.

These signed graphs that are represented by a subset of D_n , like generalized line graphs, also have a finite forbidden subgraph characterization.




Figure 8. Additional minimal forbidden subgraphs for $\mathcal{D}_{\infty}^{\pm}$ (up to switching equivalence).

Theorem 3.11 (Chawathe and Vijayakumar [8]). *The minimal forbidden subgraphs for the family $\mathcal{D}_{\infty}^{\pm}$ of signed graphs that are represented by a subset of $\mathcal{D}_n$ are listed in Figures 5 and 8 up to switching equivalence.*

Next, generalizing Lemma 2.14, we carry out the following computation.

Lemma 3.12. *For every minimal forbidden subgraph $F^{\pm}$ for the family $\mathcal{D}_{\infty}^{\pm}$, and every nonempty signed vertex subset $A^{\pm}$ of $F^{\pm}$, the path extensions and the clique extensions of $F^{\pm}$ satisfy*

$$\lim_{\ell \rightarrow \infty} \lambda_1(F^{\pm}, A^{\pm}, \ell) \leq -\lambda^* \quad \text{and} \quad \lim_{m \rightarrow \infty} \lambda_1(F^{\pm}, A^{\pm}, K_m) < -\lambda^*.$$

Moreover, the equality holds in the first inequality if and only if $F^{\pm} = G_4$ and $A^{\pm} \in \{\{3^{\pm}\}, \{4^{\pm}\}\}$.

We prove Lemma 3.12 under computer assistance in Appendix A. Lastly, the proof of Lemma 2.15 can be taken almost verbatim to show the following generalization.

Lemma 3.13. *Suppose that $A^{\pm}$ is a nonempty vertex subset of a signed graph $F^{\pm}$ and $\lambda \geq 2$. If the path extensions of $F^{\pm}$ satisfy*

$$\lim_{\ell \rightarrow \infty} \lambda_1(F^{\pm}, A^{\pm}, \ell) < -\lambda,$$

then there exists $m \in \mathbb{N}^+$ such that the path-clique extensions of $F^{\pm}$ satisfy

$$\lambda_1(F^{\pm}, A^{\pm}, \ell, K_m) < -\lambda \text{ for every } \ell \in \mathbb{N}.$$

We are ready to prove Theorem 3.9.

Proof of Theorem 3.9. Suppose that $\lambda \in [2, \lambda^*)$. Let $\mathcal{F}^{\pm}$ denote the set of minimal forbidden subgraphs for the family $\mathcal{D}_{\infty}^{\pm}$. According to Theorem 3.11, the family $\mathcal{F}^{\pm}$ is finite. Combining Lemmas 3.12 and 3.13, we choose $\ell, m \in \mathbb{N}^+$ such that for every $F^{\pm} \in \mathcal{F}^{\pm}$, the signed extension family $\mathcal{X}^{\pm}(F^{\pm}, \ell, m)$ is




Figure 9. A valid 3-coloring of a signed graph.

disjoint from $\mathcal{G}^\pm(\lambda)$. We know that $S_5 \notin \mathcal{G}^\pm(\lambda)$ and $-K_4 \notin \mathcal{G}^\pm(\lambda)$. In particular, no graph in $\mathcal{G}^\pm(\lambda)$ contains a subgraph that is switching equivalent to any member in the following family

$$\{S_5, -K_4\} \cup \bigcup_{F^\pm \in \mathcal{F}^\pm} \mathcal{X}^\pm(F^\pm, \ell, m). \quad (15)$$

Using Lemma 3.5, we obtain $N \in \mathbb{N}$ such that for every connected signed graph $G^\pm$ with more than N vertices, if no member in (15) is switching equivalent to a subgraph of $G^\pm$, then neither is any $F^\pm \in \mathcal{F}^\pm$, and so $G^\pm$ is represented by a subset of D_n and is clearly in $\mathcal{G}^\pm(2)$. This implies that every connected graph in $\mathcal{G}^\pm(\lambda) \setminus \mathcal{G}^\pm(2)$ has at most N vertices. $\square$

3.3. Proof of Theorem 1.5 for $\lambda \geq \lambda^*$

Proof of Theorem 1.5 for $\lambda \geq \lambda^$.* If $\mathcal{F}^\pm$ is a finite forbidden subgraph characterization of $\mathcal{G}^\pm(\lambda)$, then the all-positive signed graphs in $\mathcal{F}^\pm$ would form a finite forbidden subgraph characterization of $\mathcal{G}(\lambda)$, which contradicts Theorem 1.3 for $\lambda \geq \lambda^*$. $\square$

4. Spherical two-distance sets

We introduce the definition of chromatic number for signed graphs.

Definition 4.1 (Chromatic number). A *valid p -coloring* of a signed graph $G^\pm$ is a coloring of the vertices using p colors such that the endpoints of every negative edge receive different colors, and the endpoints of every positive edge receive identical colors. (See Figure 9 for an example.) The *chromatic number* $\chi(G^\pm)$ of a signed graph $G^\pm$ is the smallest p for which $G^\pm$ has a valid p -coloring. If $G^\pm$ does not have a valid p -coloring for any p , we write $\chi(G^\pm) = \infty$.

Recall from Section 1 the following spectral graph theoretic quantity,

$$k_p(\lambda) = \inf \left\{ \frac{|G^\pm|}{\text{mult}(\lambda, G^\pm)} : \chi(G^\pm) \leq p \text{ and } \lambda^1(G^\pm) = \lambda \right\}. \quad (16)$$

We say that $k_p(\lambda)$ is *achievable* if it is finite and the infimum can be attained. The spectral graph theoretic quantity $k_p(\lambda)$ can be seen as a generalization of the *spectral radius order* $k(\lambda)$ defined by

$$k(\lambda) = \min \{|G| : \lambda^1(G) = \lambda\}.$$

Indeed, $k_1(\lambda) = k_2(\lambda) = k(\lambda)$. However, the behavior of $k_p(\lambda)$ is far more mysterious when $p \geq 3$ (see the remark after [22, Definition 1.10] for more details). The technique developed in the current paper enables us to certify values of $k_p(\lambda)$ whenever $\lambda < \lambda^*$.

Theorem 4.2. For every $p \geq 3$ and $\lambda < \lambda^*$, there exists $N \in \mathbb{N}$ such that

$$k_p(\lambda) = \min \left\{ \frac{|G^\pm|}{\text{mult}(\lambda, G^\pm)} : |G^\pm| \leq N, \chi(G^\pm) \leq p \text{ and } \lambda^1(G^\pm) = \lambda \right\},$$

and in particular, $k_p(\lambda)$ is achievable whenever it is finite. Moreover, for $\lambda = 2$, there exists $p_0 \geq 3$ such that $k_p(2) = p^2/(p-1)^2$ for all $p \geq p_0$.

We will return to the proof of Theorem 4.2 towards the end of the subsection. The next result constructs large spherical codes with two fixed angles.

Proposition 4.3 (Proposition 2.2 of Jiang et al. [22]). Fix $-1 \leq \beta < 0 \leq \alpha < 1$. Then $N_{\alpha,\beta}(d) \geq d$ for every $d \in \mathbb{N}^+$. Moreover if $k_p(\lambda) < \infty$, where $\lambda = (1-\alpha)/(\alpha-\beta)$ and $p = \lfloor -\alpha/\beta \rfloor + 1$, then

$$N_{\alpha,\beta}(d) \geq \begin{cases} \frac{k_p(\lambda)d}{k_p(\lambda)-1} - O_{\alpha,\beta}(1) & \text{if } k_p(\lambda) \text{ is achievable,} \\ \frac{k_p(\lambda)d}{k_p(\lambda)-1} - o(d) & \text{otherwise.} \end{cases}$$

Conjecture 1.8 asserts that the constructions in Proposition 4.3 are optimal up to an error sublinear in d . A framework to bound $N_{\alpha,\beta}(d)$ from above was formulated in [22].

Definition 4.4 (Definition 5.2 of Jiang et al. [22]). Given $p \in \mathbb{N}^+$ and a family $\mathcal{H}$ of signed graphs, let $M_{p,\mathcal{H}}(\lambda, N)$ be the maximum possible value of $\text{mult}(\lambda, G^\pm)$ over all signed graphs $G^\pm$ on at most N vertices that do not contain any member of $\mathcal{H}$ as a subgraph and satisfy $\chi(G^\pm) \leq p$ and $\lambda^{p+1}(G^\pm) \leq \lambda$.

Theorem 4.5 (Theorem 5.3 of Jiang et al. [22]). Fix $-1 \leq \beta < 0 \leq \alpha < 1$. Set $\lambda = (1-\alpha)/(\alpha-\beta)$ and $p = \lfloor -\alpha/\beta \rfloor + 1$. Let $\mathcal{H}$ be a finite family of signed graphs with $\lambda^1(H^\pm) > \lambda$ for each $H^\pm \in \mathcal{H}$. Then

$$N_{\alpha,\beta}(d) \leq d + M_{p,\mathcal{H}}(\lambda, N_{\alpha,\beta}(d)) + O_{\alpha,\beta,\mathcal{H}}(1).$$

A proper choice of the finite family $\mathcal{H}$ in the last theorem establishes Conjecture 1.8 for $\lambda < \lambda^*$.

Corollary 4.6. Fix $-1 \leq \beta < 0 \leq \alpha < 1$. Set $\lambda = (1-\alpha)/(\alpha-\beta)$ and $p = \lfloor -\alpha/\beta \rfloor + 1$. If $\lambda < \lambda^*$, then

$$N_{\alpha,\beta}(d) = \begin{cases} \frac{k_p(\lambda)d}{k_p(\lambda)-1} + O_{\alpha,\beta}(1) & \text{if } k_p(\lambda) < \infty, \\ d + O_{\alpha,\beta}(1) & \text{otherwise.} \end{cases}$$

Proof. According to Corollary 1.6, we can take a finite forbidden subgraph characterization, denoted $\mathcal{H}$, of the family $\mathcal{G}^\mp(\lambda)$ of signed graphs with largest eigenvalue at most λ . Thus $M_{p,\mathcal{H}}(\lambda, N)$ is the maximum possible value of $\text{mult}(\lambda, G^\pm)$ over all signed graphs $G^\pm$ on at most N vertices that satisfy $\chi(G^\pm) \leq p$ and $\lambda^1(G^\pm) \leq \lambda$. Note that $M_{p,\mathcal{H}}(\lambda, N)$ is a nondecreasing function of N . We break the rest of the proof into two cases.

Case 1: $M_{p,\mathcal{H}}(\lambda, N) = 0$ for every $N \in \mathbb{N}^+$. In other words, there is no signed graph $G^\pm$ that satisfy $\chi(G^\pm) \leq p$ and $\lambda^1(G^\pm) = \lambda$. Thus this case corresponds to the case where $k_p(\lambda) = \infty$. According to Proposition 4.3 and Theorem 4.5, we have $N_{\alpha,\beta}(d) = d + O_{\alpha,\beta}(1)$.

Case 2: $M_{p,\mathcal{H}}(\lambda, N) > 0$ for every $N \geq d_0$. This case corresponds to the case where $k_p(\lambda) < \infty$. Suppose that $d \geq d_0$. Proposition 4.3 says that $N_{\alpha,\beta}(d) \geq d$, and so $M_{p,\mathcal{H}}(\lambda, N_{\alpha,\beta}(d)) > 0$. Let $G^\pm$ be a signed graph with at most $N_{\alpha,\beta}(d)$ vertices that satisfy $\chi(G^\pm) \leq p$, $\lambda^1(G^\pm) \leq \lambda$ and $\text{mult}(\lambda, G^\pm) = M_{p,\mathcal{H}}(\lambda, N_{\alpha,\beta}(d))$. Thus

$$k_p(\lambda) \leq \frac{|G^\pm|}{\text{mult}(\lambda, G^\pm)} \leq \frac{N_{\alpha,\beta}(d)}{M_{p,\mathcal{H}}(\lambda, N_{\alpha,\beta}(d))}.$$

Theorem 4.5 then implies that

$$N_{\alpha,\beta}(d) \leq \frac{k_p(\lambda)d}{k_p(\lambda) - 1} + O_{\alpha,\beta}(1),$$

which, in view of Theorem 4.2 and Proposition 4.3, gives $N_{\alpha,\beta}(d) = k_p(\lambda)d/(k_p(\lambda)-1) + O_{\alpha,\beta}(1)$. $\square$

Lastly, we come back to the achievability of the spectral graph theoretic quantity $k_p(\lambda)$. Actually we may assume in the definition (16) of $k_p(\lambda)$ that $G^\pm$ is connected by passing to a suitable connected component of $G^\pm$ in case $G^\pm$ is not connected. Furthermore, as we shall mainly work with the signed graph that reverses all the edge signs of $G^\pm$ in (16), we redefine $k_p(\lambda)$ as follows:

$$k_p(\lambda) = \inf \left\{ \frac{|G^\pm|}{\text{mult}(-\lambda, G^\pm)} : G^\pm \text{ is connected, } \chi(-G^\pm) \leq p, \text{ and } \lambda_1(G^\pm) = -\lambda \right\},$$

where $-G^\pm$ reverses all the edge signs of $G^\pm$.

We break the proof of Theorem 4.2 into three cases $\lambda < 2$, $\lambda \in (2, \lambda^*)$, and $\lambda = 2$.

Proof of Theorem 4.2 for $\lambda < 2$. We obtain $\ell \in \mathbb{N}^+$ from Lemma 2.3(p) such that $P_{\ell+1} \notin \mathcal{G}^\pm(\lambda)$. Applying Lemma 3.5 to $F^\pm = K_1$ (the 1-vertex graph), $k_1 = 4$, $k_2 = 3$ and $m = 2p$, we obtain $N \in \mathbb{N}$ such that for every connected signed graph $G^\pm$, if $G^\pm$ does not contain any subgraph that is switching equivalent to any member in $\{S_4, -K_3\} \cup \mathcal{X}^\pm(K_1, \ell, 2p)$, then $G^\pm$ contains at most N vertices.

It suffices to show that for every connected signed graph $G^\pm$ with more than N vertices, either $\chi(-G^\pm) > p$ or $G^\pm \notin \mathcal{G}^\pm(\lambda)$. By our choice of N , the signed graph $G^\pm$ contains a subgraph that is switching equivalent to a member in $\{S_4, -K_3\} \cup \mathcal{X}^\pm(K_1, \ell, 2p)$. We break the rest of the proof into two cases.

Case 1: $G^\pm$ contains a subgraph that is switching equivalent to S_4 , $-K_3$, or a path extension in $\mathcal{X}^\pm(K_1, \ell, 2p)$. Since $\lambda_1(S_4) = \lambda_1(-K_3) = -2$ and every path extension in $\mathcal{X}^\pm(K_1, \ell, 2p)$ is switching equivalent to $P_{\ell+1}$, the signed graph $G^\pm$ is not in $\mathcal{G}^\pm(\lambda)$.

Case 2: $G^\pm$ contains a subgraph that is switching equivalent to a path-clique extension or a clique extension in $\mathcal{X}^\pm(K_1, \ell, 2p)$. Observe that every path-clique extension and every clique extension in $\mathcal{X}^\pm(K_1, \ell, 2p)$ contains a subgraph that is switching equivalent to K_{2p+1} , and moreover every signed graph that is switching equivalent to K_{2p+1} always contains K_{p+1} as a subgraph. Thus the signed graph $G^\pm$ contains K_{p+1} as a subgraph. Therefore $-G^\pm$ contains $-K_{p+1}$ as a subgraph, and so $\chi(-G^\pm) \geq p+1$. $\square$

The proof of Theorem 4.2 for $\lambda \in (2, \lambda^*)$ follows immediately from Theorem 3.9.

Proof of Theorem 4.2 for $\lambda \in (2, \lambda^)$.* Theorem 3.9 implies that there exists $N \in \mathbb{N}$ such that every connected signed graph $G^\pm$ with $\lambda_1(G^\pm) = -\lambda$ has at most N vertices. $\square$

The proof of Theorem 4.2 for $\lambda = 2$ is trickier. We need the following generalization of Theorem 2.17. This generalization was explicitly recognized by Chawathe and Vijayakumar [8, Theorem 1.1], who attributed the result to Witt [33], and referred the reader to [7] for a combinatorial proof.

Theorem 4.7 (Witt [33] and Cameron et al. [7]). *For every connected signed graph $G^\pm$, the smallest eigenvalue of $G^\pm$ is at least -2 if and only if $G^\pm$ is represented by a subset of D_n or E_8 .*

Recall that $\mathcal{D}_\infty^\pm$ denotes the family of signed graphs that are represented by a subset of D_n for some $n \in \mathbb{N}^+$, where

$$D_n := \{\varepsilon_1 e_i + \varepsilon_2 e_j : \varepsilon_1, \varepsilon_2 = \pm 1, 1 \leq i < j \leq n\}.$$

We shall focus on the signed graphs in $\mathcal{D}_\infty^\pm$. To that end, we generalize bidirected graphs to multigraphs as follows.




Figure 10. A proper 3-edge-coloring of a bidirected multigraph.

Definition 4.8 (Bidirected multigraph). A *bidirected multigraph* is a multigraph (allowing parallel edges, but no loops) in which each vertex-edge incidence (v, e) has a positive or negative sign, denoted $\sigma(v, e) \in \{\pm 1\}$, such that every pair of parallel edges e_1 and e_2 satisfies the following condition: for one of the shared endpoints, say v_1 , the incidences (v_1, e_1) and (v_1, e_2) have the same sign, while for the other shared endpoint v_2 , the incidences (v_2, e_1) and (v_2, e_2) have opposite signs. In particular, there are at most two edges between any two vertices in a bidirected multigraph.

As we transfer relevant properties of a signed graph $G^\pm \in \mathcal{D}_\infty^\pm$ to a bidirected multigraph $\vec{H}$, a vertex coloring of $G^\pm$ will correspond to an edge coloring of $\vec{H}$.

Definition 4.9 (Intersecting edges and proper edge colorings). As opposed to parallel edges, we say that two edges of a bidirected multigraph *intersect* at a vertex v if v is the only vertex they have in common. A *proper p -edge-coloring* of a bidirected multigraph $\vec{H}$ is a coloring of the edges using p colors such that for each pair of edges e_1, e_2 that intersect at one vertex, say v , the edges e_1 and e_2 receive different colors when the incidences (v, e_1) and (v, e_2) have the same sign, whereas e_1 and e_2 receive identical colors when (v, e_1) and (v, e_2) have opposite signs. See Figure 10 for an example.

Definition 4.10 (Associated vectors and uniform vertices). Suppose that $v_1, \dots, v_n$ are the vertices of a bidirected multigraph $\vec{H}$. For every edge $e \in E(\vec{H})$ with endpoints v_i and v_j , define the *associated vector* of the edge e by

$$\vec{e} := \sigma(v_i, e)\mathbf{e}_i + \sigma(v_j, e)\mathbf{e}_j \in D_n.$$

We say that a vertex v of $\vec{H}$ is *uniform* if the sign $\sigma(v, e)$ is the same for every edge e incident to v .

Proposition 4.11. *There is a one-to-one correspondence between signed graphs $G^\pm$ in $\mathcal{D}_\infty^\pm$ and bidirected multigraphs $\vec{H}$ without isolated vertices such that the following properties hold.*

- (1) *The underlying graph of $G^\pm$ is the line graph of $\vec{H}$, in particular, the number of vertices of $G^\pm$ is equal to the number of edges of $\vec{H}$.*
- (2) *The signed graph $G^\pm$ is connected if and only if $\vec{H}$ is connected except when $\vec{H}$ consists of a single pair of parallel edges.*
- (3) *The signed graph $-G^\pm$ has a valid p -coloring if and only if $\vec{H}$ has a proper p -edge-coloring.*
- (4) *The multiplicity $\text{mult}(-2, G^\pm)$ is equal to $m - \text{rank } V$, where m is the number of edges of $\vec{H}$ and $V = \{\vec{e} \in D_n : e \in E(\vec{H})\}$.*
- (5) *The signed graph $G^\pm$ is all-positive if and only if every vertex of $\vec{H}$ is uniform.*

Proof. Suppose that $G^\pm$ is represented by $V \subseteq D_n$. Construct a bidirected multigraph $\vec{H}$ with vertices $v_1, \dots, v_n$ and incidence signing σ as follows. For each vector in V of the form $\varepsilon_1\mathbf{e}_i + \varepsilon_2\mathbf{e}_j$ with $i < j$, we put an edge e connecting v_i and v_j , and we assign $\sigma(v_i, e) = \varepsilon_1$ and $\sigma(v_j, e) = \varepsilon_2$. For every pair of parallel edges e_1 and e_2 , which connect v_i and v_j , since $\sigma(v_i, e_1)\mathbf{e}_i + \sigma(v_j, e_1)\mathbf{e}_j$ and $\sigma(v_i, e_2)\mathbf{e}_i + \sigma(v_j, e_2)\mathbf{e}_j$ are two distinct vectors from V , their inner product satisfies

$$\{0, \pm 2\} \ni \sigma(v_i, e_1)\sigma(v_i, e_2) + \sigma(v_j, e_1)\sigma(v_j, e_2) = A_{G^\pm}(e_1, e_2) \in \{0, \pm 1\},$$

and so their inner product must be 0. Therefore $\vec{H}$ is a bona fide bidirected multigraph. We may remove all the isolated vertices (if any) from $\vec{H}$.

Conversely one can read off the vectors in V from $\vec{H}$ – for every edge e , include the associated vector $\vec{e} \in D_n$ in V . Furthermore one can reconstruct $G^\pm$ from $\vec{H}$ as follows. We identify the vertex set of

$G^\pm$ with the edge set of $\vec{H}$, and for each pair of distinct vertices e_1 and e_2 of $G^\pm$, they are adjacent in $G^\pm$ if and only if e_1 and e_2 intersect at one vertex, say v , in $\vec{H}$, and the sign of the edge e_1e_2 in $G^\pm$ is $\sigma(v, e_1)\sigma(v, e_2)$. One can then check that the reconstructed $G^\pm$ is indeed represented by V .

Finally, it is routine to transfer properties back and forth between $G^\pm$ and $\vec{H}$. $\square$

Now we are ready to strengthen the last case of Theorem 4.2.

Lemma 4.12. *For every $p \geq 3$, every connected signed graph $G^\pm$ in $\mathcal{D}_\infty^\pm$ with $\chi(-G^\pm) \leq p$ and $\lambda_1(G^\pm) = -2$ satisfies*

$$\frac{|G^\pm|}{\text{mult}(-2, G^\pm)} \geq \left(\frac{p}{p-1}\right)^2,$$

and equality holds if and only if $G^\pm$ is the all-positive line graph of the complete bipartite graph $K_{p,p}$.

Proof. According to Proposition 4.11, it suffices to show that for every connected bidirected multigraph $\vec{H}$ with n vertices and m edges, if $\vec{H}$ has a proper p -edge-coloring, then

$$\frac{m}{m - \text{rank } V} \geq \left(\frac{p}{p-1}\right)^2 \text{ or equivalently } \frac{m}{\text{rank } V} \leq \frac{p^2}{2p-1}, \text{ where } V := \{\vec{e} \in D_n : e \in E(\vec{H})\},$$

and equality holds if and only if $\vec{H}$ is the complete bipartite graph $K_{p,p}$ with uniform vertices.

Suppose that $c: E(\vec{H}) \rightarrow \{c_1, \dots, c_p\}$ is a proper p -edge-coloring of $\vec{H}$. One can deduce from Definition 4.9 the following properties.

- (a) For two edges e_1 and e_2 that intersect at a vertex v , if $\sigma(v, e_1) = \sigma(v, e_2)$, then $c(e_1) \neq c(e_2)$.
- (b) For two edges e_1 and e_2 that intersect at a vertex v , if $\sigma(v, e_1) \neq \sigma(v, e_2)$, then $c(e_1) = c(e_2)$, and any other edge incident to v has to be parallel to either e_1 or e_2 .

We break the rest of the proof on the upper bound of $m/\text{rank } V$ into two cases.

Case 1: $\vec{H}$ has no parallel edges. We claim that the maximum degree of $\vec{H}$ is at most p . Suppose on the contrary that k edges $e_1, \dots, e_k$ pairwise intersect at a vertex v , where $k > p$. Because $k > p \geq 3$, the contrapositive of (b) implies that $\sigma(v, e_i)$ is the same for $i \in \{1, \dots, k\}$. Moreover (a) says that $c(e_1), \dots, c(e_k)$ are distinct, which contradicts the assumption that c is a proper p -edge-coloring. In particular, the claim implies that

$$m/n \leq p/2. \tag{17}$$

Since $\vec{H}$ is connected, one can check that the set of vectors $\{\vec{e} \in D_n : e \in E(T)\}$, where T is a spanning tree of $\vec{H}$, is linearly independent. Therefore $\text{rank } V \in \{n-1, n\}$. We may assume that

$$4 \leq n \leq 2p \quad \text{and} \quad \text{rank } V = n-1.$$

Indeed, in the case where $n \leq 3$, we have

$$\frac{m}{\text{rank } V} \leq \frac{m}{n-1} \leq \frac{\binom{n}{2}}{n-1} = \frac{n}{2} \leq \frac{3}{2} < \frac{9}{5} \leq \frac{p^2}{2p-1},$$

in the case where $n > 2p$, we have

$$\frac{m}{\text{rank } V} \leq \frac{m}{n-1} \stackrel{(17)}{\leq} \frac{p/2}{1-1/n} < \frac{p/2}{1-1/(2p)} = \frac{p^2}{2p-1}.$$

and in the case where $\text{rank } V = n$, we also have

$$\frac{m}{\text{rank } V} = \frac{m}{n} \stackrel{(17)}{\leq} \frac{p}{2} < \frac{p^2}{2p-1},$$

Let U be the vertex set of $\vec{H}$, and define the vertex subset

$$U_0 := \{u \in U : u \text{ is not uniform}\}.$$

We claim that $\vec{H}[U \setminus U_0]$ is bipartite. To specify its bipartition, let $\mathbf{x} : U \rightarrow \mathbb{R}$ be a nonzero vector in the orthogonal complement of V . For every edge $e \in E(\vec{H})$ with endpoints u_1 and u_2 , since $\mathbf{x}$ is orthogonal to the vector $\vec{e}$ associated to e , we know that

$$\sigma(u_1, e)\mathbf{x}(u_1) + \sigma(u_2, e)\mathbf{x}(u_2) = 0, \quad (18)$$

hence $|\mathbf{x}(u_1)| = |\mathbf{x}(u_2)|$. Because $\vec{H}$ is connected, we deduce that $|\mathbf{x}(u)|$ is constant for every $u \in U$, and in particular, $\mathbf{x}(u) \neq 0$ for every $u \in U$. Since every vertex in $U \setminus U_0$ is uniform, we partition $U \setminus U_0$ as follows:

$$U_1 := \{u \in U \setminus U_0 : \sigma(u, e)\mathbf{x}(u) > 0 \text{ for every edge } e \text{ incident to } u\},$$

$$U_2 := \{u \in U \setminus U_0 : \sigma(u, e)\mathbf{x}(u) < 0 \text{ for every edge } e \text{ incident to } u\}.$$

In view of (18), we deduce that $\vec{H}[U \setminus U_0]$ is a bipartite graph with parts U_1 and U_2 , and so the number of edges in $\vec{H}[U \setminus U_0]$ is at most $|U_1||U_2|$. According to (b), every vertex in U_0 has degree at most 2. Thus, we can estimate m using $n_0 := |U_0|$ as follows:

$$m \leq |U_1||U_2| + 2|U_0| \leq \left(\frac{n-n_0}{2}\right)^2 + 2n_0 \leq \max\left(\frac{n^2}{4}, 2n-3\right),$$

where the last inequality assumes that $0 \leq n_0 \leq n-2$. In the corner case where $n_0 \geq n-1$, we can estimate m by

$$2m \leq (n-1)(|U_1| + |U_2|) + 2|U_0| = (n-1)(n-n_0) + 2n_0 \leq 3(n-1),$$

which implies $m \leq 3(n-1)/2 < 2n-3$. Therefore, we always have

$$\frac{m}{\text{rank } V} = \frac{m}{n-1} \leq \max\left(\frac{n^2}{4(n-1)}, \frac{2n-3}{n-1}\right) = \begin{cases} \frac{2n-3}{n-1} & \text{if } n < 6, \\ \frac{n^2}{4(n-1)} & \text{if } n \geq 6, \end{cases}$$

which, as a function of n , increases on $(1, \infty)$. As $n \leq 2p$ and $p \geq 3$, the function reaches its maximum $p^2/(2p-1)$ at $n = 2p$, and so $m/\text{rank } V \leq p^2/(2p-1)$. It is easy to see that equality holds if and only if $\vec{H}$ is the complete bipartite graph $K_{p,p}$ with uniform vertices.

Case 2: $\vec{H}$ has parallel edges. Let e_1 and e_2 be a pair of parallel edges. Since $\vec{H}$ is connected, there is a spanning tree T that contains e_1 . One can check that the set of vectors $\{\vec{e} \in D_n : e \in E(T) \cup \{e_2\}\}$ is linearly independent. Therefore

$$\text{rank } V = n.$$

We next use the discharging method to bound the average degree $2m/n$ of $\vec{H}$. Each vertex v starts with the charge $d(v)$, that is, the degree of v . We claim that for every vertex v with $d(v) > p$, exactly one of the following two situations happens.

- (I) The vertex v is uniform, that is, the sign $\sigma(v, e)$ is the same for every edge e incident to v .
 (II) The edges incident to v are precisely two pairs (e_1, e'_1) and (e_2, e'_2) of parallel edges satisfying $\sigma(v, e_1) \neq \sigma(v, e_2)$, and in particular $d(v) = 4$ and $p = 3$.

Indeed, suppose that $d(v) > p$, and suppose that situation (I) does not happen, that is, there exist two edges e_1 and e_2 incident to v such that $\sigma(v, e_1) \neq \sigma(v, e_2)$. Because $d(v) > p \geq 3$, we may further assume that e_1 and e_2 intersect at v by replacing e_1 or e_2 with another edge incident to v in case e_1 and e_2 are parallel. Because $d(v) \geq 4$, according to (b), situation (II) must happen.

To describe the discharging rules, we say that two edges e_1 and e_2 are *twin* if they are parallel and $c(e_1) = c(e_2)$, and for every vertex v with $d(v) > p$, we call v a *type I* vertex when (I) happens, and a *type II* vertex when (II) happens.


- o The first discharging rule sends 1 from every type I vertex v to each of its neighbors v' that are connected to v through twin edges.
- o The second discharging rule sends $1/2$ from every type II vertex v to each of its two neighbors.

One can further deduce from Definition 4.9 the following property of twin edges.

- (c) For twin edges e_1 and e_2 incident to a vertex v , if $\sigma(v, e_1) \neq \sigma(v, e_2)$, then e_1 and e_2 are the only two edges incident to v .

We first consider the charge of each vertex after the first discharging rule is applied. Suppose that v is a type I vertex, and v' is an arbitrary vertex that is connected to v through a pair of parallel edges e_1 and e_2 . Since $\sigma(v, e_1) = \sigma(v, e_2)$, it must be the case that $\sigma(v', e_1) \neq \sigma(v', e_2)$, and so v' is not a type I vertex. Thus v does not receive any charge from its neighbors. According to (a), among all the edges incident to v , only the parallel ones can receive identical colors. Thus after v sends out charges to its neighbors, its charge decreases exactly to the number of distinct colors assigned to the edges incident to v , and so the charge of v is at most p now. Suppose in addition that e_1 and e_2 are twin edges, or in other words, v' receives 1 from v . According to (c), the only edges incident to v' are e_1 and e_2 , and so the charge of v' is at most 3, which is at most p , now.

We then consider the charge of each vertex after the second discharging rule is applied. Notice that the second discharging rule only kicks in when $p = 3$. Suppose that v is a type II vertex, and two pairs (e_1, e'_1) and (e_2, e'_2) of parallel edges connect v respectively to v_1 and v_2 such that $\sigma(v, e_1) \neq \sigma(v, e_2)$. Without loss of generality, we may assume that $\sigma(v, e_1) = +1$ and $\sigma(v, e_2) = -1$. We have the four possibilities for $\sigma(v, e'_1)$ and $\sigma(v, e'_2)$. Once $\sigma(v, e'_1)$ and $\sigma(v, e'_2)$ are chosen, we can determine, according to (a) and (b), which edges incident to v receive identical or distinct colors. We summarize the four possibilities in the following figure. For each possibility, it is easy to check that v did not receive any charge from v_1 or v_2 when the first discharging rule was applied. Therefore the final charge of v is 3, which is equal to p .



We are left to decide the final charge of v_1 and v_2 . Observe that for $i \in \{1, 2\}$ there are, *au fond*, only three possible ways, as shown below, to color e_i and e'_i and to assign $\sigma(v_i, e_i)$ and $\sigma(v_i, e'_i)$. For the first possibility, according to (c), the only edges incident to v_i are e_i and e'_i , and so the final charge of v_i is $5/2$, which is less than p . For the other two possibilities, the contrapositive of (b) implies that v_i must be uniform, and so its charge was at most p after the first discharging rule was applied. Assume for the sake of contradiction that v_i is connected to another type II vertex w through edges f_i and f'_i . Since v_i is uniform, in view of the three possibilities above, we know that $c(e_i) \neq c(e'_i)$ and $c(f_i) \neq c(f'_i)$.

In view of (a), we know that $\{c(e_i), c(e'_i)\} \cap \{c(f_i), c(f'_i)\} = \emptyset$. Thus the four edges e_i, e'_i, f_i and f'_i receive distinct colors under c , which contradicts the assumption that $p = 3$. Therefore v is the only type II neighbor of v_i , and the final charge of v_i is at most $p + 1/2$.



Since the discharging rules preserve the total charge, the average degree $2m/n$ is at most the maximum final charge, which is at most $p + 1/2$. Finally, we obtain that

$$\frac{m}{\text{rank } V} = \frac{m}{n} \leq \frac{p + 1/2}{2} < \frac{p^2}{2p - 1}. \quad \square$$

Proof of Theorem 4.2 for $\lambda = 2$. From Theorem 4.7 and Lemma 4.12, we know that $k_p(2)$ is the smaller of the two quantities $p^2/(p - 1)^2$ and

$$k_p^* := \inf \left\{ \frac{|G^\pm|}{\text{mult}(-2, G^\pm)} : G^\pm \text{ is represented by a subset of } E_8 \text{ and } \chi(-G^\pm) \leq p \right\}.$$

Because there are only finitely many signed graphs that are represented by a subset of E_8 , the infimum in the definition of k_p^* is in fact a minimum, hence $k_p(2)$ is achievable. Moreover $k_p^* \geq k^*$, where

$$k^* := \min \left\{ \frac{|G^\pm|}{\text{mult}(-2, G^\pm)} : G^\pm \text{ is represented by a subset of } E_8 \right\}. \quad (19)$$

Since $k^* > 1$, there exists $p_0 \geq 3$ such that $p^2/(p - 1)^2 \leq k^* \leq k_p^*$ for every $p \geq p_0$. $\square$

5. Concluding remarks

Extending Smith's classification [29] of connected graphs with spectral radius at most 2, Cvetković, Doob and Gutman [9, Theorem 3.8] characterized the connected graphs with spectral radius in $(2, \lambda')$, which were later explicitly classified by Brouwer and Neumaier [4]. Recall that $\lambda' = \sqrt{2} + \sqrt{5} \approx 2.05817$. Retrospectively, perhaps both Theorem 1.2 and the fact that λ' is the smallest $\lambda \in \mathbb{R}$ such that the set of graph spectral radii is dense in (λ, ∞) (cf. [16, 28]) indicated that it is possible to classify connected graphs with spectral radius at most λ' .

A similar line of research was carried out for signed graphs. McKee and Smyth [24] determined all the connected signed graphs with spectral radius at most 2 – they are switching equivalent to subgraphs of S_{14} , S_{16} , and T_{2n} for $n \geq 3$ in Figure 11. When investigating Lehmer's Mahler measure problem, McKee and Smyth [25, Theorem 3] further determined the 17 connected signed graphs with spectral radius in $(2, 2.019)$. Belardo, Cioabă, Koolen and Wang [3, Problem 3.11] raised the question on the classification of connected signed graphs with spectral radius at most λ' . This question was very recently resolved by Wang, Dong, Hou and Li [32].

With regard to smallest eigenvalues, we would like to extend Theorem 2.17 of Cameron et al. [7] beyond $\mathcal{G}(2)$.

Problem 5.1. Classify all the connected graphs with smallest eigenvalue in $(-\lambda^*, -2)$. In particular, classify such graphs that have sufficiently many vertices.

It is worth mentioning that Bussemaker and Neumaier [6, Theorem 2.5] showed that $E_{2,6}$ defined in Proposition 2.1 is the only connected graph with smallest eigenvalue in $[-\lambda^1(E_{2,6}), -2)$, where




Figure 11. Maximal connected signed graphs with spectral radius at most 2 up to switching equivalence. The number of vertices in T_{2n} is $2n$.

$\lambda^1(E_{2,6}) \approx 2.00659$. Very recently, Acharya and Jiang [1] provided a complete solution to Problem 5.1 – there are 794 infinite families of graphs and 4,752 exceptional graphs.

We can ask the same question for signed graphs, extending Theorem 4.7 beyond $\mathcal{G}^\pm(2)$.

Problem 5.2. Classify all the connected signed graphs with smallest eigenvalue in $(-\lambda^*, -2)$. In particular, classify such signed graphs that have sufficiently many vertices.

Turning to spherical two-distance sets with two fixed angles, it is plausible to establish more instances of Conjecture 1.8 by taking advantage of $\chi(G^\pm)$ in Definition 4.4. Denote by $\mathcal{G}_p^\pm(\lambda)$ the family of signed graphs $G^\pm$ with $\chi(G^\pm) \leq p$ and $\lambda^1(G^\pm) \leq \lambda$. Observe that $\mathcal{G}_p^\pm(\lambda)$, just like $\mathcal{G}^\pm(\lambda)$, is still closed under taking subgraphs. We raise the following question, whose solution could possibly establish more instances of Conjecture 1.8.

Problem 5.3. For every $p \in \mathbb{N}^+$, determine the set of $\lambda \in \mathbb{R}$ for which $\mathcal{G}_p^\pm(\lambda)$ has a finite forbidden subgraph characterization.

Notice that $\mathcal{G}_1^\pm(\lambda)$ consists of unsigned graphs only, and $\mathcal{G}_2^\pm(\lambda)$ consists of signed graphs that are switching equivalent to unsigned graphs. Thus Theorem 1.2 essentially answers Problem 5.3 when $p \in \{1, 2\}$.

Finally, we point out that the proof of Theorem 4.2 for $\lambda = 2$ actually shows that $k_p(2) = p^2/(p-1)^2$ for every $p \geq 1/(1-1/\sqrt{k^*})$, where k^* is defined as in (19). We compute $k^* = 15/14$ hence $1/(1-1/\sqrt{k^*}) \approx 29.49$. Indeed, Stanić noted in [30] that, up to switching equivalence, there is a unique maximal signed graph, denoted $M_8^\pm$, that is represented by a subset of E_8 , and moreover, the spectrum of $M_8^\pm$ is $\{28^8, (-2)^{112}\}$. By the Cauchy interlacing theorem, every signed graph $G^\pm$ that is represented by a subset of E_8 satisfies $\lambda^1(G^\pm) \leq 28$, and thus

$$0 = \sum_i \lambda_i(G^\pm) \leq (-2) \text{mult}(-2, G^\pm) + 28(|G^\pm| - \text{mult}(-2, G^\pm)),$$

which implies that $|G^\pm|/\text{mult}(-2, G^\pm) \geq 15/14$. Since equality holds when $G^\pm = M_8^\pm$, we obtain $k^* = 15/14$. We leave the determination of $k_p(2)$ for small p as an open problem.

Problem 5.4. For every $p \in \{3, 4, \dots, 29\}$, determine the value of $k_p(2)$.

A. Computer-assisted proofs

Notice that the path extensions $(G_4, \{3^\pm\}, \ell)$ and $(G_4, \{4^\pm\}, \ell)$ are switching equivalent to $E_{2,\ell+3}$ (see Figure 2), and so Proposition 2.1 implies that equality holds for $F = G_4$ and $A \in \{\{3\}, \{4\}\}$ in Lemma 2.14, and for $F^\pm = G_4$ and $A \in \{\{3^\pm\}, \{4^\pm\}\}$ in Lemma 3.12.

The termination of a program that solves the following computational problem is a proof of the strict inequalities in Lemmas 2.14 and 3.12. In fact, we strengthen these strict inequalities by replacing

Table 1. Input

G1	0312142334
G2	02030412142334
G3	0102030534
G4	0102030405
G5	01020304051234
G6	01020512152345
G7	01020304052345
G8	0312131415233445
G9	0102030405233445
G10	020304051215232534
G11	01020305121545
G12	0102051215233445
G13	0205121523253435
G14	0312131415233445
G15	0205121523253545
G16	01020304051215232534
G17	010203051215232534
G18	020304051215232534
G19	010203051215232534
G20	010203040512152345
G21	010203040512152325
G22	020512152324253545
G23	02121314152324343545
G24	01020304051215233445
G25	01030412131415233445
G26	01020304051213141534
G27	0102030412131415233445
G28	0102030405121314152345
G29	010203040512131415233445
G30	010203040512152324253545
G31	01020304051215232425343545
S32	011213-23
S33	02031213-23
S34	02031213-0123
S35	0102031213-23
S36	0112152345-34
S37	0304121523-01
S38	030412152345-10
S39	030412152334-01
S40	010203052345-34
S41	03041215233445-01
S42	01020512152345-34
S43	01020405121523-34
S44	02030515232545-34
S45	0203121415232545-34
S46	0203051215232545-34
S47	010203051214152345-34
S48	010203051215232545-34
S49	01020305121415232545-34

$\lambda^* \approx 2.0198$ with $101/50 = 2.02$. Note that $-101/50$ is not an algebraic integer, and hence it cannot be an eigenvalue of any signed graph.

Input. The first line of the input gives the number N of minimal forbidden subgraphs for $\mathcal{D}_\infty^\pm$ (up to switching equivalence). Each of the N lines that follow represents a signed graph in Figures 5 and 8 by two strings. The first string is the label of the signed graph. The second string is of the form $u[1]u[2] \dots u[2e-1]u[2e]$ possibly followed by $-v[1]v[2] \dots v[2f-1]v[2f]$, which lists the positive edges $u[1]u[2], \dots, u[2e-1]u[2e]$ and the negative edges $v[1]v[2], \dots, v[2f-1]v[2f]$.

Output. For each of the N signed graphs $F^\pm$, output one line containing $x \ y \ z$, where x is the label of $F^\pm$, y and z are both -1 if $\lambda_1(F^\pm) < -101/50$ already, otherwise y is, in all but one case, the minimum ℓ such that $\lambda_1(F^\pm, A^\pm, \ell) < -101/50$ for every nonempty signed vertex subset $A^\pm$ of $F^\pm$, and z is the minimum m such that $\lambda_1(F^\pm, A^\pm, K_m) < -101/50$ for every nonempty signed vertex subset $A^\pm$ of $F^\pm$. In the exceptional case where the label of $F^\pm$ is G4, y is defined similarly but $A^\pm$ cannot be $\{3^\pm\}$ or $\{4^\pm\}$, and an extra $*$ is appended to y .

Our implementation is straightforward. We iterate through the minimal forbidden subgraphs $F^\pm$ for $\mathcal{D}_\infty^\pm$. In each iteration, we compute y and z as follows. We first check whether $A_{F^\pm} + (101/50)I$ is positive definite – if not we output $-1 \ -1$ for $y \ z$ and continue to the next iteration. We test whether a matrix is positive definite by checking whether its leading principal minors are all positive. If the smallest eigenvalue of $F^\pm$ turns out to be more than $-101/50$, we then go through all possible nonempty signed vertex subsets $A^\pm$ of $F^\pm$. For each $A^\pm$, we increase ℓ from 0 until the determinant of $A_{(F^\pm, A^\pm, \ell)} + (101/50)I$ is negative, and we increase m from 1 until the determinant of $A_{(F^\pm, A^\pm, K_m)} + (101/50)I$ is negative. We record the largest ℓ and m when we go through $A^\pm$, and output them as y and z . In the exceptional case where the label of $F^\pm$ is G4, we skip the computation of ℓ when $A^\pm$ is $\{3^\pm\}$ or $\{4^\pm\}$. We avoid floating-point errors by representing every number as a rational number.

The actual code, written in Ruby, is available as the ancillary file `maximum_extensions.rb` in the arXiv version of this paper. As our code halts, we obtain a proof of Lemmas 2.14 and 3.12. We provide the input in Table 1 for the convenience of anyone who wants to program independently, and we provide in addition the output in Table 2 for cross-check.

Table 2. Output

G1 -1 -1	G2 -1 -1	G3 -1 -1	G4 8* 7	G5 -1 -1	G6 -1 -1
G7 6 5	G8 11 7	G9 -1 -1	G10 7 6	G11 7 6	G12 6 6
G13 4 4	G14 5 5	G15 -1 -1	G16 5 5	G17 -1 -1	G18 4 5
G19 6 5	G20 4 5	G21 5 5	G22 -1 -1	G23 6 5	G24 4 4
G25 4 4	G26 -1 -1	G27 4 4	G28 4 5	G29 4 4	G30 -1 -1
G31 4 4	S32 -1 -1	S33 -1 -1	S34 -1 -1	S35 -1 -1	S36 -1 -1
S37 6 5	S38 4 4	S39 6 5	S40 5 5	S41 4 4	S42 -1 -1
S43 4 5	S44 5 5	S45 5 5	S46 4 4	S47 4 4	S48 4 4
S49 4 4					

Acknowledgments. We thank Sebastian Cioabă for sending us a copy of [18], and Jing Zhang for pointing us to [13]. We are very grateful to Zhuang Xiong and two anonymous referees for their valuable feedback on the earlier versions of the paper. One referee noted a gap in the proof of a byproduct result of Theorem 1.5 on symmetric hollow integer matrices in an earlier version; that result was removed from the present manuscript and later proved in [19].

Competing interest. The authors have no competing interests to declare.

Financial support. ZJ was supported in part by an AMS Simons Travel Grant, and by U.S. taxpayers through NSF grant DMS-2127650. AP was supported in part by U.S. taxpayers through NSF grant DMS-2349045.

References

- [1] H. Acharya and Z. Jiang, ‘Beyond the classification theorem of Cameron, Goethals, Seidel, and Shult’, Preprint, 2024, [arXiv:2404.13136](https://arxiv.org/abs/2404.13136) [math.CO].
- [2] I. Balla, F. Dräxler, P. Keevash and B. Sudakov, ‘Equiangular lines and spherical codes in Euclidean space’, *Invent. Math.* **211**(1) (2018), 179–212, [arXiv:1606.06620](https://arxiv.org/abs/1606.06620) [math.CO].
- [3] F. Belardo, S. M. Cioabă, J. Koolen and J. Wang, ‘Open problems in the spectral theory of signed graphs’, *Art Discrete Appl. Math.* **1**(2) (2018), Paper No. 2.10, 23, [arXiv:1907.04349](https://arxiv.org/abs/1907.04349) [math.CO].
- [4] A. E. Brouwer and A. Neumaier, ‘The graphs with spectral radius between 2 and $\sqrt{2 + \sqrt{5}}$ ’, *Linear Algebra Appl.* **114** / 115 (1989), 273–276.
- [5] B. Bukh, ‘Bounds on equiangular lines and on related spherical codes’, *SIAM J. Discrete Math.* **30**(1) (2016), 549–554, [arXiv:1508.00136](https://arxiv.org/abs/1508.00136) [math.CO].
- [6] F. C. Bussemaker and A. Neumaier, ‘Exceptional graphs with smallest eigenvalue -2 and related problems’, *Math. Comp.* **59**(200) (1992), 583–608, with microfiche supplement.
- [7] P. J. Cameron, J.-M. Goethals, J. J. Seidel and E. E. Shult, ‘Line graphs, root systems, and elliptic geometry’, *J. Algebra* **43**(1) (1976), 305–327.
- [8] P. D. Chawathe and G. R. Vijayakumar, ‘A characterization of signed graphs represented by root system D_∞ ’, *European J. Combin.* **11**(6) (1990), 523–533.
- [9] D. Cvetković, M. Doob and I. Gutman, ‘On graphs whose spectral radius does not exceed $(2 + \sqrt{5})^{1/2}$ ’, *Ars Combin.* **14** (1982), 225–239.
- [10] D. Cvetković, M. Doob and S. Simić, ‘Some results on generalized line graphs’, *C. R. Math. Rep. Acad. Sci. Canada* **2**(3) (1980), 147–150.
- [11] D. Cvetković, M. Doob and S. Simić, ‘Generalized line graphs’, *J. Graph Theory* **5**(4) (1981), 385–399.
- [12] D. Cvetković, P. Rowlinson and S. Simić, *Spectral Generalizations of Line Graphs: On Graphs with Least Eigenvalue -2* , London Mathematical Society Lecture Note Series, vol. 314 (Cambridge Univ. Press, Cambridge, 2004).
- [13] L. E. Dickson, ‘Finiteness of the odd perfect and primitive abundant numbers with n distinct prime factors’, *Amer. J. Math.* **35**(4) (1913), 413–422.
- [14] M. Doob, ‘The limit points of eigenvalues of graphs’, *Linear Algebra Appl.* **114** / 115 (1989), 659–662.
- [15] G. Greaves, J. Koolen, A. Munemasa, Y. Sano and T. Taniguchi, ‘Edge-signed graphs with smallest eigenvalue greater than -2 ’, *J. Combin. Theory Ser. B* **110** (2015), 90–111, [arXiv:1309.5178](https://arxiv.org/abs/1309.5178) [math.CO].
- [16] A. J. Hoffman, ‘On limit points of spectral radii of non-negative symmetric integral matrices’, in *Graph Theory and Applications* (Proc. Conf., Western Michigan Univ., Kalamazoo, Mich., 1972; dedicated to the memory of J. W. T. Youngs), Lecture Notes in Mathematics, vol. 303 (Springer, 1972), 165–172.
- [17] A. J. Hoffman, ‘On graphs whose least eigenvalue exceeds $-1 - \sqrt{2}$ ’, *Linear Algebra Appl.* **16**(2) (1977), 153–165.
- [18] A. J. Hoffman, ‘On limit points of the least eigenvalue of a graph’, *Ars Combin.* **3** (1977), 3–14.

- [19] Z. Jiang, 'On symmetric hollow integer matrices with eigenvalues bounded from below', *Linear Algebra Appl.* **709** (2025), 233–240, [arXiv:2408.16860](#) [math.CO].
- [20] Z. Jiang and A. Polyanskii, 'Forbidden subgraphs for graphs of bounded spectral radius, with applications to equiangular lines', *Israel J. Math.* **236**(1) (2020), 393–421, [arXiv:1708.02317](#) [math.CO].
- [21] Z. Jiang, J. Tidor, Y. Yao, S. Zhang and Y. Zhao, 'Equiangular lines with a fixed angle', *Ann. Math. (2)* **194**(3) (2021), 729–743, [arXiv:1907.12466](#) [math.CO].
- [22] Z. Jiang, J. Tidor, Y. Yao, S. Zhang and Y. Zhao, 'Spherical two-distance sets and eigenvalues of signed graphs', *Combinatorica* **43**(2) (2023), 203–232, [arXiv:2006.06633](#) [math.CO].
- [23] V. Kumar, S. B. Rao and N. M. Singhi, 'Graphs with eigenvalues at least -2 ', *Linear Algebra Appl.* **46** (1982), 27–42.
- [24] J. McKee and C. Smyth, 'Integer symmetric matrices having all their eigenvalues in the interval $[-2, 2]$ ', *J. Algebra* **317**(1) (2007), 260–290, [arXiv:0705.3599](#) [math.CO].
- [25] J. McKee and C. Smyth, 'Integer symmetric matrices of small spectral radius and small Mahler measure', *Int. Math. Res. Not. IMRN* **2012**(1) (2012), 102–136, [arXiv:0907.0371](#) [math.NT].
- [26] S. B. Rao, N. M. Singhi and K. S. Vijayan, 'The minimal forbidden subgraphs for generalized line-graphs', in *Combinatorics and Graph Theory (Calcutta, 1980)*, Lecture Notes in Mathematics, vol. 885 (Springer, Berlin-New York, 1981), 459–472.
- [27] A. C. M. van Rooij and H. S. Wilf, 'The interchange graph of a finite graph', *Acta Math. Acad. Sci. Hungar.* **16** (1965), 263–269.
- [28] J. B. Shearer, 'On the distribution of the maximum eigenvalue of graphs', *Linear Algebra Appl.* **114** / 115 (1989), 17–20.
- [29] J. H. Smith, 'Some properties of the spectrum of a graph', in *Combinatorial Structures and Their Applications* (Proc. Calgary Internat. Conf., Calgary, Alta., 1969), Gordon and Breach, New York (1970), 403–406.
- [30] Z. Stanić, 'Notes on exceptional signed graphs', *Ars Math. Contemp.* **18**(1) (2020), 105–115.
- [31] G. R. Vijayakumar, 'Signed graphs represented by D_{∞} ', *European J. Combin.* **8**(1) (1987), 103–112.
- [32] D. Wang, W. Dong, Y. Hou and D. Li, 'On signed graphs whose spectral radius does not exceed $\sqrt{2} + \sqrt{5}$ ', *Discrete Math.* **346**(6) (2023), 113358, [arXiv:2203.01530](#) [math.CO].
- [33] E. Witt, 'Spiegelungsgruppen und Aufzählung halbeinfacher Liescher Ringe', *Abh. Math. Sem. Hansischen Univ.* **14** (1941), 289–322.
- [34] T. Zaslavsky, 'Matrices in the theory of signed simple graphs', in *Advances in Discrete Mathematics and Applications: Mysore, 2008*, Ramanujan Mathematical Society Lecture Notes Series, vol. 13 (Ramanujan Math. Soc., Mysore, 2010), 207–229, [arXiv:1303.3083](#) [math.CO].