



RESEARCH ARTICLE

Two uniqueness results in the inverse boundary value problem for the weighted p -Laplace equation

Catalin Carstea¹ and Ali Feizmohammadi²

¹Department of Applied Mathematics, National Yang Ming Chiao Tung University, Hsinchu 300, Taiwan, ROC;
E-mail: catalin.carstea@gmail.com (corresponding author).

²Department of Mathematics, University of Toronto, Mississauga, ON L5L 1C6, Canada;
E-mail: ali.feizmohammadi@utoronto.ca.

Received: 9 November 2024; Revised: 26 June 2025; Accepted: 26 June 2025

2020 Mathematical Subject Classification: Primary – 35R30; Secondary – 35J62, 35J70, 35J92

Abstract

In this paper we prove a general uniqueness result in the inverse boundary value problem for the weighted p -Laplace equation in the plane, with smooth weights. We also prove a uniqueness result in dimension 3 and higher, for real analytic weights that are subject to a smallness condition on one of their directional derivatives. Both results are obtained by linearizing the equation at a solution without critical points. This unknown solution is then recovered, together with the unknown weight.

1. Introduction

Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be a compact connected set with nonempty interior and a smooth boundary, let $\gamma \in C^\infty(\Omega)$ be a positive function¹, and finally let $p \in (1, 2) \cup (2, \infty)$. We consider the boundary value problem

$$\begin{cases} \nabla \cdot (\gamma |\nabla u|^{p-2} \nabla u) = 0, \\ u|_{\partial\Omega} = f, \end{cases} \quad (1)$$

where f, u are real-valued functions. Equation (1) is known as the *weighted p -Laplace equation* and it is a quasilinear, degenerate elliptic equation. The forward problem for this equation is well studied and we have

Theorem 1.1 (e.g., [33, Theorem 1]). *Let $f \in C^{1,\alpha}(\partial\Omega)$ for some $\alpha \in (0, 1]$. There exist $\beta \in (0, 1)$ and $C(\|f\|_{C^{1,\alpha}(\partial\Omega)}) > 0$ nondecreasing such that equation (1) has a unique weak solution $u \in C^{1,\beta}(\Omega)$ and*

$$\|u\|_{C^{1,\beta}(\Omega)} \leq C(\|f\|_{C^{1,\alpha}(\partial\Omega)}). \quad (2)$$

It is therefore possible to define the Dirichlet-to-Neumann map associated to (1) by

$$\Lambda_\gamma(f) = \left(\gamma |\nabla u|^{p-2} \partial_\nu u \right) \Big|_{\partial\Omega}, \quad \forall f \in C^{1,\alpha}(\partial\Omega), \quad (3)$$

where u is the unique solution to (1) and ν is the exterior normal unit vector on $\partial\Omega$.

¹Note on notation: since we choose Ω to denote a closed set, $C^\infty(\Omega)$ is the space that other works might be denoted by $C^\infty(\bar{\Omega})$, i.e. the space of restrictions of $C^\infty(\mathbb{R}^n)$ functions to $\Omega = \bar{\Omega}$.

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

In [7], Calderón proposed the following question/inverse problem: Can an elliptic coefficient γ be recovered from the Dirichlet-to-Neumann map associated to the equation $\nabla \cdot (\gamma \nabla u) = 0$? A positive answer for general smooth coefficients γ was first provided in [42] in dimension 3 or higher and by [35] in the plane. In the intervening decades, similar questions for other equations have been investigated in a large number of papers. It is beyond the purposes of our work to give a full account of the existing inverse problems literature. We will reference below those works that are most closely related to our own, in terms of subject matter or technique.

In this paper we are interested in the inverse problem of recovering the a priori unknown coefficient γ in (1), given the knowledge of Λ_γ . This is a natural analogue of the original problem of Calderón. We will prove two results. The first is the following general uniqueness result in the plane.

Theorem 1.2. *Let $n = 2$ and let $\gamma, \tilde{\gamma} \in C^\infty(\Omega)$ be strictly positive functions. If $\Lambda_\gamma = \Lambda_{\tilde{\gamma}}$, then $\gamma = \tilde{\gamma}$.*

In dimensions 3 and higher we prove the following uniqueness result for weights belonging to $C^\omega(\Omega)$, the space of real-analytic functions on Ω .

Theorem 1.3. *Let $n \geq 3$ and let $\zeta \in \mathbb{R}^n$ be a unit vector. Suppose there exists a point $z \in \partial\Omega$ in a neighborhood of which $\partial\Omega$ is flat. There exists $\mu > 0$, depending only on Ω and n , such that if $\gamma, \tilde{\gamma} \in C^\omega(\Omega)$ are strictly positive functions with $\|\zeta \cdot \nabla \gamma\|_{C^{0,\alpha}(\Omega)}, \|\zeta \cdot \nabla \tilde{\gamma}\|_{C^{0,\alpha}(\Omega)} < \mu$, then $\Lambda_\gamma = \Lambda_{\tilde{\gamma}}$ implies $\gamma = \tilde{\gamma}$.*

The study of inverse problems for nonlinear equations is not new, but in recent years there has been a considerable increase in the interest for this topic. As examples, we can cite the papers [17], [21], [23], [24], [28], [29], [31], [30], [40] on semilinear equations, and [10], [9], [11], [14], [15], [8], [16], [20], [22], [25], [34], [37], [38], [39], [41] on quasilinear equations. By and large, all these works rely on a so-called *second/higher linearization* method. These ideas go back to [21], where what one may call a first-order linearization method was used. As commonly employed, the method consists of using Dirichlet data that depends on a small (or large) parameter ϵ , typically of the form $\epsilon\phi$ (or $\lambda + \epsilon\phi$, with λ a constant, if constants are a solution to the linear part of the equation). One then uses the asymptotic expansion of the Dirichlet-to-Neumann map in terms of the parameter ϵ to obtain information about the coefficients of the equation. Sometimes this is presented as differentiating the equation with respect to the small parameter, then setting it to zero.

Our paper is not the first to take up the inverse boundary value problem for the weighted p -Laplacian. The works [6], [3], [4], [5], [19], [27], [36] all address aspects of the same problem. We note that a uniqueness result without additional constraints, such as monotonicity, has not yet been previously derived for the weighted p -Laplacian. Also, past boundary determination results have only yielded $\gamma|_{\partial\Omega}$ and $\partial_\nu \gamma|_{\partial\Omega}$, but not the rest of the derivatives of γ on the boundary (see [36], [3]). For other degenerate equations, the only known results are those of [12], [13], where general uniqueness results are derived for the coefficients of porous medium equations.

The approach to the proofs of Theorems 1.2 and 1.3 also makes use of a linearization method. In equation (1) we use Dirichlet data of the form $f = \phi_0 + \epsilon\phi$, with ϵ a small parameter. Let u_ϵ be the corresponding solution and, assuming that we are justified in taking the derivative, let $\dot{u} = \frac{d}{d\epsilon} u_\epsilon|_{\epsilon=0}$. Further assuming we can differentiate the equation, it is not hard to see that $\dot{u}$ should satisfy the anisotropic linear equation

$$\begin{cases} \nabla \cdot (A \nabla \dot{u}) = 0, \\ \dot{u}|_{\partial\Omega} = \phi, \end{cases} \quad (4)$$

where A is the matrix with the u_0 -dependent coefficients

$$A_{jk} = \gamma |\nabla u_0|^{p-2} \left(\delta_{jk} + (p-2) \frac{\partial_j u_0 \partial_k u_0}{|\nabla u_0|^2} \right). \quad (5)$$

The Dirichlet-to-Neumann map Λ_γ determines the Dirichlet-to-Neumann map Λ_A for the equation (4).

In order to use already established results for the determination of the coefficient matrix A , we need it to be elliptic. Indeed, even the differentiability of u_ϵ w.r.t. ϵ is in question unless that is the case. We then see that the unknown u_0 must be guaranteed to have no critical points. In dimension 2, by results of Alessandrini and Sigalotti in [2], we can guarantee the absence of critical points by choosing Dirichlet data that has single local minimum and maximum points on $\partial\Omega$. In dimension 3 and higher something like this is unlikely to hold, as even for linear elliptic equations it is known that for each Dirichlet data there is an open set of smooth coefficients that produce solutions with critical points (see [1]). We can show, however, that, for coefficients γ that vary slowly in one direction, there exists explicit Dirichlet data for which no critical points appear.

We can also point out here a simple corollary of our linearization result (Proposition 2.3 below), for weights that are constant in one direction.

Corollary 1.1. *Let $n \geq 3$ and $\zeta \in \mathbb{R}^n$ be a unit vector. If $\gamma, \tilde{\gamma} \in C^\infty(\Omega)$ are such that $\zeta \cdot \nabla \gamma = \zeta \cdot \nabla \tilde{\gamma} = 0$ and $\Lambda_\gamma = \Lambda_{\tilde{\gamma}}$, then $\gamma = \tilde{\gamma}$.*

Proof. In this case $u_0 = \zeta \cdot x$ is a solution of (1), with either weight. Then

$$A_{jk} = \gamma(\delta_{jk} + (p-2)\zeta_j\zeta_k), \quad \tilde{A}_{jk} = \tilde{\gamma}(\delta_{jk} + (p-2)\zeta_j\zeta_k). \quad (6)$$

After a rescaling in the ζ direction, the linearized problem reduces to the classical Calderón problem with isotropic conductivities. $\square$

The linearization procedure is detailed in section 2. In section 3 we give a proof of Theorem 1.2. By the well-known result [35] of Nachman, we have uniqueness for the coefficient matrix A , up to diffeomorphism invariance. Making use of the particular structure of A , we then succeed in showing that the diffeomorphism relating A and $\tilde{A}$ must be trivial and that $\gamma = \tilde{\gamma}$. In section 4 we give the proof of Theorem 1.3. Our approach is to use boundary determination results for equation (4) to obtain the values of all tangential directions of A on the boundary, together with all their normal direction derivatives. From this information we are then able to inductively show uniqueness for the values of all the normal direction derivatives $\partial_\nu^k u_0|_{\partial\Omega}$, $\partial_\nu^k \gamma|_{\partial\Omega}$, $k = 0, 1, 2, \dots$. Since here γ is assumed to be a real-analytic function, this is enough to recover it on Ω .

2. Linearizing the p -Laplace equation

For each $\xi \in \mathbb{R}^n \setminus \{0\}$ let

$$J_j(\xi) = |\xi|^{p-2}\xi_j, \quad j = 1, \dots, n. \quad (7)$$

Then

$$\frac{\partial}{\partial \xi_k} J_j(\xi) = |\xi|^{p-2} \left(\delta_{jk} + (p-2) \frac{\xi_j \xi_k}{|\xi|^2} \right). \quad (8)$$

In what follows, we will repeatedly use Taylor's formula

$$J_j(\zeta) = J_j(\xi) + \sum_{k=1}^n (\zeta_k - \xi_k) \int_0^1 \partial_{\xi_k} J(\xi + t(\zeta - \xi)) dt. \quad (9)$$

We plan to linearize equation (1) near some solution u_0 , whose boundary data $u_0|_{\partial\Omega} = \phi_0$ is known. As will become apparent below, we can only perform the linearization if u_0 does not have any critical points in Ω . In dimension two plenty of such solutions exist, thanks to the following proposition due to Alessandrini and Sigalotti.

Proposition 2.1 (see [2, Theorem 5.1]). *If $n = 2$ there exists boundary data $\phi_0 \in C^\infty(\partial\Omega)$ independent of γ such that the corresponding solution u_0 of (1) is in $C^\infty(\Omega)$ and $|\nabla u_0(x)| > 0$ for any $x \in \Omega$.*

In higher dimensions, even for a linear elliptic equation with unknown coefficients it is impossible to guarantee the absence of critical points (see [1]). We can still show the existence of such a solution provided the weight γ is sufficiently close to a constant.

Proposition 2.2. *Let $\zeta \in \mathbb{R}^n$ be a unit vector. There exists $\mu > 0$ so that if $\|\zeta \cdot \nabla \gamma\|_{C^{0,\alpha}(\Omega)} < \mu$, then there exists $u_0 \in C^\infty(\Omega)$ which solves (1) with boundary data $\phi_0 = \zeta \cdot x$, and is such that $|\nabla u_0(x)| > 0$ for any $x \in \Omega$.*

Proof. Without loss of generality, we assume that $\zeta = (1, 0, \dots, 0)$. We make the Ansatz

$$u_0(x) = x_1 + R, \quad R|_{\partial\Omega} = 0. \quad (10)$$

By (9) we have

$$\sum_k B_{jk}(\nabla R) \partial_k R = \gamma J_j(\nabla u_0) - \gamma \delta_{1j}, \quad (11)$$

$$B_{jk}(\xi) = \gamma \int_0^1 |e_1 + t\xi|^{p-2} \left(\delta_{jk} + (p-2) \frac{(\delta_{1j} + t\xi_j)(\delta_{1k} + t\xi_k)}{|e_1 + t\xi|^2} \right) dt. \quad (12)$$

Taking the divergence of the above we get

$$\begin{cases} \nabla \cdot (B(\nabla R) \nabla R) = -\partial_1 \gamma, \\ R|_{\partial\Omega} = 0. \end{cases} \quad (13)$$

Let $V \in C^{2,\alpha}(\Omega)$ be such that $\|V\|_{C^{2,\alpha}(\Omega)} < 1/2$ and define the map $T(V) = U$, where U is the solution to

$$\begin{cases} \nabla \cdot (B(\nabla V) \nabla U) = -\partial_1 \gamma, \\ U|_{\partial\Omega} = 0. \end{cases} \quad (14)$$

Since $B(\nabla V) \in C^{1,\alpha}(\Omega)$ are uniformly elliptic coefficients, it follows that a unique solution $U \in C^{2,\alpha}(\Omega)$ exists (see [18, Theorem 6.14]). Furthermore, by [18, Theorem 6.6] we have

$$\|U\|_{C^{2,\alpha}(\Omega)} \leq C \|\partial_1 \gamma\|_{C^{0,\alpha}(\Omega)}. \quad (15)$$

If the right hand side is less than $1/2$, by Schauder's fixed point theorem (see [18, Theorem 11.1]) it follows that T has a fixed point on the ball of radius $1/2$ in $C^{2,\alpha}(\Omega)$. By uniqueness of solutions for (1), this must be R and we conclude that

$$\|\nabla R\|_{L^\infty(\Omega)} \leq \frac{1}{2}, \quad (16)$$

so

$$|\nabla u_0(x)| > \frac{1}{2}, \quad \forall x \in \Omega. \quad (17)$$

Note that the nonvanishing of the gradient ∇u_0 makes the equation satisfied by u_0 elliptic, so the smoothness of u_0 follows. $\square$

In what follows we will assume that u_0 is as in the preceding two propositions. Let A be the matrix with coefficients

$$A_{jk} = \gamma |\nabla u_0|^{p-2} \left(\delta_{jk} + (p-2) \frac{\partial_j u_0 \partial_k u_0}{|\nabla u_0|^2} \right). \quad (18)$$

Proposition 2.3. *Under the assumptions of either Proposition 2.1 or Proposition 2.2, we have that the Dirichlet-to-Neumann map Λ_γ for the weighted p -Laplace equation (1) determines the Dirichlet-to-Neumann map Λ_A for the linear equation $\nabla \cdot (A \nabla u) = 0$, on the same domain Ω .*

Proof. For $\phi \in C^\infty(\partial\Omega)$, and $\epsilon \in \mathbb{R}$ small, let u_ϵ be the solution of

$$\begin{cases} \nabla \cdot (\gamma |\nabla u_\epsilon|^{p-2} \nabla u_\epsilon) = 0, \\ u_\epsilon|_{\partial\Omega} = u_0|_{\partial\Omega} + \epsilon \phi. \end{cases} \quad (19)$$

We make the Ansatz

$$u_\epsilon = u_0 + R_\epsilon. \quad (20)$$

By Theorem 1.1 and the theorem of Arzelà-Ascoli, it follows that (on a subsequence) we have that $R_\epsilon \rightarrow R_0$ in $C^1(\Omega)$. Since then $u_0 + R_0$ would be a weak solution of the same boundary value problem u_0 satisfies, it follows that $R_\epsilon \rightarrow 0$ in $C^1(\Omega)$. Since the limit is the same for every subsequence, it follows that in fact we do not need to pass to a subsequence. This is easily seen as follows: suppose there is a subsequence of R_{ϵ_k} of R_ϵ so that $\liminf_{\epsilon_k \rightarrow 0^+} \|R_{\epsilon_k} - R_0\|_{C^1(\Omega)} > 0$; however, the argument above also shows that R_{ϵ_k} has a subsequence which converges to R_0 in $C^1(\Omega)$, which is a contradiction.

By Taylor's formula we have that

$$\sum_k \partial_k R_\epsilon \int_0^1 \partial_{\xi_k} J_j(\nabla u_0 + t \nabla R_\epsilon) dt = J_j(\nabla u_\epsilon) - J_j(\nabla u_0). \quad (21)$$

Let

$$A_{jk}^\epsilon = \gamma \int_0^1 \partial_{\xi_k} J_j(\nabla u_0 + t \nabla R_\epsilon) dt. \quad (22)$$

Since $R_\epsilon \rightarrow 0$ in C^1 , it follows that $|\nabla u_0 + t \nabla R_\epsilon|$ is uniformly bounded and uniformly bounded away from zero. This implies that A_{jk}^ϵ is a set of elliptic coefficients, with ellipticity bounds independent of ϵ . Taking gradients in (21) we get that R_ϵ satisfies

$$\begin{cases} \nabla \cdot (A^\epsilon \nabla R_\epsilon) = 0, \\ R_\epsilon|_{\partial\Omega} = \epsilon \phi, \end{cases} \quad (23)$$

It follows that

$$\|R_\epsilon\|_{C^{1,\beta}(\Omega)} \leq C\epsilon. \quad (24)$$

We can again invoke the theorem of Arzelà-Ascoli to conclude that there must exist $\dot{u} \in C^1(\Omega)$ such that $\epsilon^{-1} R_\epsilon \rightarrow \dot{u}$ in $C^1(\Omega)$. Taking the limit in (23) we see that $\dot{u}$ must be a weak solution of

$$\begin{cases} \nabla \cdot (A \nabla \dot{u}) = 0, \\ \dot{u}|_{\partial\Omega} = \phi, \end{cases} \quad (25)$$

Returning to (21), dividing by ϵ and taking the limit $\epsilon \rightarrow 0$, we have that

$$\begin{aligned} \nu \cdot A \nabla \dot{u} &= \lim_{\epsilon \rightarrow 0} \gamma \frac{\nu \cdot J(\nabla u_\epsilon) - \nu \cdot J(\nabla u_0)}{\epsilon} \\ &= \lim_{\epsilon \rightarrow 0} \frac{\Lambda_\gamma(u_0|_{\partial\Omega} + \epsilon \phi) - \Lambda_\gamma(u_0|_{\partial\Omega})}{\epsilon}. \end{aligned} \quad (26)$$

We see then that the Neumann data for the equation (25) is determined by the map Λ_γ . $\square$

3. Proof of Theorem 1.2

Suppose $n = 2$ and we have $\gamma, \tilde{\gamma}$ as above such that $\Lambda_\gamma = \Lambda_{\tilde{\gamma}}$. We use notation such as $u_\epsilon, \tilde{u}_\epsilon$ to denote the corresponding solutions to (19), etc. Observe that one consequence of the identity of the DN maps is

$$\int_{\Omega} \gamma |\nabla u_0|^p \, dx = \int_{\Omega} \tilde{\gamma} |\nabla \tilde{u}_0|^p \, dx. \quad (27)$$

By Proposition 2.3 we have $\Lambda_A = \Lambda_{\tilde{A}}$. From [35, Theorem 2] it follows that there must exist a diffeomorphism $\Phi = (\Phi^1, \Phi^2) : \Omega \rightarrow \Omega$ such that $\Phi|_{\partial\Omega} = Id$ and

$$\tilde{A}(x) = \frac{1}{|D\Phi|} (D\Phi)^T A D\Phi \circ \Phi^{-1}(x). \quad (28)$$

Here $D\Phi$ is the matrix with coefficients $(D\Phi)_{jk} = \partial_j \Phi^k$.

Note that

$$\begin{aligned} & \det \left(\delta_{jk} + (p-2) \frac{\partial_j u_0 \partial_k u_0}{|\nabla u_0|^2} \right) \\ &= \left(1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) \left(1 + (p-2) \frac{(\partial_2 u_0)^2}{|\nabla u_0|^2} \right) \\ &\quad - (p-2)^2 \frac{(\partial_1 u_0 \partial_2 u_0)^2}{|\nabla u_0|^4} = p-1. \end{aligned} \quad (29)$$

Taking determinants on both sides of (28) we obtain

$$\left(\tilde{\gamma} |\nabla \tilde{u}_0|^{p-2} \right)^2 = \left(\gamma |\nabla u_0|^{p-2} \right)^2 |D\Phi|^{-2} |D\Phi|^2 \circ \Phi^{-1}, \quad (30)$$

so

$$\tilde{\gamma} |\nabla \tilde{u}_0|^{p-2} = \gamma |\nabla u_0|^{p-2} \circ \Phi^{-1}. \quad (31)$$

Another consequence of $\Lambda_A = \Lambda_{\tilde{A}}$ is that for each ϕ we have $\dot{\tilde{u}} = \dot{u} \circ \Phi^{-1}$. Incidentally, for $\phi = u_0|_{\partial\Omega}$ the solution to the linear equation is $\dot{u} = u_0$. Therefore

$$\tilde{u}_0 = u_0 \circ \Phi^{-1}. \quad (32)$$

It then follows that

$$\nabla u_0 = D\Phi(\nabla \tilde{u}_0 \circ \Phi), \quad (33)$$

which we can use in (28), together with (31), to get

$$\begin{aligned} \tilde{A} &= \tilde{\gamma} |\nabla \tilde{u}_0|^{p-2} \frac{1}{|D\Phi| \circ \Phi^{-1}} \left([(D\Phi)^T D\Phi] \circ \Phi^{-1} + (p-2) \frac{|\nabla \tilde{u}_0|^2}{|\nabla u_0|^2 \circ \Phi^{-1}} \right. \\ &\quad \left. \times [(D\Phi)^T D\Phi] \circ \Phi^{-1} \frac{\nabla \tilde{u}_0 \otimes \nabla \tilde{u}_0}{|\nabla \tilde{u}_0|^2} [(D\Phi)^T D\Phi] \circ \Phi^{-1} \right). \end{aligned} \quad (34)$$

Let

$$F = \left[\frac{(D\Phi)^T D\Phi}{|D\Phi|} \right] \circ \Phi^{-1}, \quad P = \frac{\nabla \tilde{u}_0 \otimes \nabla \tilde{u}_0}{|\nabla \tilde{u}_0|^2}, \quad \alpha = |D\Phi| \circ \Phi^{-1} \frac{|\nabla \tilde{u}_0|^2}{|\nabla u_0|^2 \circ \Phi^{-1}}. \quad (35)$$

Note that F, P are symmetric matrices and that $P^2 = P$. Identifying P, F , and α in (34) and (18), we have

$$F + \alpha(p - 2)FPF = I + (p - 2)P. \quad (36)$$

As both $I + (p - 2)P$ and F are invertible and the left-hand side of (36) can be factored as either $F(I + (p - 2)PF)$ or $(I + (p - 2)FP)F$, it follows that both $I + \alpha(p - 2)PF$ and $I + \alpha(p - 2)FP$ are also invertible.

If we multiply (36) by P on the left, we get

$$PF + \alpha(p - 2)PFPF = (p - 1)P, \quad (37)$$

so

$$PF = (p - 1)(I + \alpha(p - 2)PF)^{-1}P, \quad (38)$$

since the inverse exists. Multiplying by P on the right and using that $P^2 = P$ yields

$$\begin{aligned} PFP &= (p - 1)(I + \alpha(p - 2)PF)^{-1}P^2 \\ &= (p - 1)(I + \alpha(p - 2)PF)^{-1}P = PF. \end{aligned} \quad (39)$$

On the other hand,

$$FP + \alpha(p - 2)FPFP = (p - 1)P, \quad (40)$$

so

$$FP = (p - 1)P(I + \alpha(p - 2)FP)^{-1}, \quad (41)$$

and therefore

$$FP = PFP = PF. \quad (42)$$

Since F and P commute, they can be simultaneously diagonalized. Since P is a rank-one projection matrix, we can write

$$F = \theta P + \eta(I - P), \quad \theta, \eta \text{ scalars.} \quad (43)$$

Multiplying (36) by $I - P$ and using the commutativity of F and P together with the identity $P(I - P) = 0$, it is easy to see that $\eta = 1$. Multiplying (36) by P and using (43) yields

$$(\theta + \alpha(p - 2)\theta^2)P = (p - 1)P \quad \Rightarrow \quad \theta + \alpha(p - 2)\theta^2 = p - 1. \quad (44)$$

On the other hand, taking the determinant of the definition of F in (35) and also in (43) gives

$$1 = \frac{|D\Phi|^2}{|D\Phi|^2} \circ \Phi^{-1} = \det F = \theta\eta = \theta. \quad (45)$$

It follows that $\theta = 1$, $F = I$, and by (44) also that $\alpha = 1$.

Suppose a nontrivial diffeomorphism such as Φ exists. Let σ be a scalar conductivity on Ω and let

$$\sigma_*(y) = \frac{\sigma}{|D\Phi|} (D\Phi)^T D\Phi \circ \Phi^{-1}(y) = \sigma \circ \Phi^{-1}(y) F(y) = \sigma \circ \Phi^{-1}(y). \quad (46)$$

This new conductivity is also scalar and gives the same DN map as σ . This violates the known uniqueness results for the Calderón problem in the plane (e.g., see [35]). So Φ must be trivial. Therefore $u_0 = \tilde{u}_0$ and $\gamma = \tilde{\gamma}$.

4. Proof of Theorem 1.3

As in the previous section, we will denote by u_0 , $\tilde{u}_0$, A , $\tilde{A}$, etc. the functions corresponding to the coefficients γ and $\tilde{\gamma}$ respectively. By Proposition 2.3 we have that $\Lambda_A = \Lambda_{\tilde{A}}$.

It is an immediate consequence of [32, Proposition 1.3] (or [26, Theorem 1.3]) that there must exist a neighborhood of U of $\partial\Omega$ and a smooth diffeomorphism $\Phi : U \cap \Omega \rightarrow U \cap \Omega$, for which we will also use the notation $\Phi = (\Phi^1, \dots, \Phi^n)$, with $\Phi|_{\partial\Omega} = Id$, and such that

$$\partial_\nu^j \tilde{A} \Big|_{\partial\Omega} = \partial_\nu^j \frac{1}{|D\Phi|} (D\Phi)^T A D\Phi \Big|_{\partial\Omega}, \quad j = 0, 1, 2, \dots \quad (47)$$

Let $z \in \partial\Omega$. Unless otherwise specified, all the following computations will be pointwise, at this point z . We wish to proceed inductively in the order of differentiation in (47).

0th order:

We have that

$$\tilde{A}(z) = \frac{1}{|D\Phi|(z)} (D\Phi)^T(z) A(z) D\Phi(z). \quad (48)$$

If τ is any unit tangent vector to $\partial\Omega$ at z , we must have that $D\Phi(z)\tau = \tau$. Since $u_0|_{\partial\Omega} = \tilde{u}_0|_{\partial\Omega}$, we also have that $\tau \cdot \nabla u_0(z) = \tau \cdot \nabla \tilde{u}_0(z)$. Therefore

$$\tau \cdot \tilde{A}(z)\tau = \tilde{\gamma}(z) |\nabla \tilde{u}_0|^{p-2}(z) \left(1 + (p-2) \frac{(\tau \cdot \nabla u_0)^2(z)}{|\nabla \tilde{u}_0|^2(z)} \right). \quad (49)$$

On the other hand, by (48) we have

$$\tau \cdot \tilde{A}(z)\tau = \frac{1}{|D\Phi|(z)} \gamma(z) |\nabla u_0|^{p-2}(z) \left(1 + (p-2) \frac{(\tau \cdot \nabla u_0)^2(z)}{|\nabla u_0|^2(z)} \right). \quad (50)$$

We can vary τ in the tangent space to the boundary at z , which is at least two-dimensional. Our plan is to use two different choices for τ .

Note that the intersection of the space of vectors that are orthogonal to $\nabla u_0(z)$ with the tangent space to the boundary at z has dimension at least $n-2$, so it cannot be trivial. By choosing $\tau \perp \nabla u_0(z)$ we can separately identify

$$\tilde{\gamma}(z) |\nabla \tilde{u}_0|^{p-2}(z) = \frac{1}{|D\Phi|(z)} \gamma(z) |\nabla u_0|^{p-2}(z). \quad (51)$$

It is, in principle, possible for the tangent space to the boundary at z to coincide with the space of vectors that are orthogonal to $\nabla u_0(z)$. Recall that on the boundary we are choosing $u_0(x) = \zeta \cdot x$, therefore $\nabla u_0(z)$ is orthogonal to the boundary at z if and only if ζ is. If that is the case, note that a unit vector $\zeta' \in \mathbb{R}^n$ that is sufficiently close, but not identical, to ζ will still satisfy the condition in the statement of Proposition 2.2. Therefore, without loss of generality, we may assume that ζ is not orthogonal to the boundary at z , and so neither is $\nabla u_0(z)$. Choosing now a tangent vector τ such that $\tau \not\perp \nabla u_0(z)$ we get

$$\frac{1}{|\nabla \tilde{u}_0|^2}(z) = \frac{1}{|\nabla u_0|^2}(z). \quad (52)$$

It follows that

$$|\nabla u_0|(z) = |\nabla \tilde{u}_0|(z), \quad (53)$$

and, as we already know that $\tau \cdot \nabla u_0(z) = \tau \cdot \nabla \tilde{u}_0(z)$ for all τ as above, we also conclude that

$$\partial_\nu u_0(z) = \partial_\nu \tilde{u}_0(z). \quad (54)$$

As $\Lambda_\gamma(u_0|_{\partial\Omega}) = \Lambda_{\tilde{\gamma}}(u_0|_{\partial\Omega})$, we get

$$\gamma(z) = \tilde{\gamma}(z). \quad (55)$$

This further implies that

$$|D\Phi|(z) = 1, \quad (56)$$

Since now $A(z) = \tilde{A}(z)$ and $D\Phi$ acts as the identity in the tangent space to $\partial\Omega$ at z , equation (48) can only hold if

$$D\Phi(z) = I. \quad (57)$$

1st order:

We have that

$$(\partial_\nu \tilde{A})(z) = \left(\partial_\nu \frac{1}{|D\Phi|} (D\Phi)^T A D\Phi \right)(z). \quad (58)$$

From this point onward, we will use the assumption that $\partial\Omega$ is flat in a neighborhood of z . For ease of computation, we will rotate our coordinates so that $\nu = e_1$ and locally $\partial\Omega \cap U \subset \{x_1 = 0\}$. We also find it notationally convenient to introduce the tangential gradient $\nabla' = \nabla - \partial_1 e_1$. As above, by possibly slightly changing the vector ζ in the statement of Proposition 2.2, we can make sure that ζ is not tangent to $\partial\Omega$ at z . Since ∇u_0 is close to ζ in L^∞ norm, we may assume, without loss of generality, that $\partial_1 u_0(z) \neq 0$.

In the previous step we have shown that $D\Phi(z) = I$. It follows that

$$\partial_j \partial_k \Phi^l(z) = 0, \quad \text{unless } j = k = 1. \quad (59)$$

Rewriting (58) with this information, we obtain that at z

$$\partial_1 \tilde{A}_{jk} = \partial_1 A_{jk} + (A_{j1} \partial_1^2 \Phi^k + A_{1k} \partial_1^2 \Phi^j) - A_{jk} \partial_1^2 \Phi^1. \quad (60)$$

In preparation for using the equations above and denoting by a_{11}, a_{jj}, a_{j1} terms made up of quantities for which uniqueness has already been shown in the previous step, that is, they depend on $\gamma|_{\partial\Omega \cap U}$, $u_0|_{\partial\Omega \cap U}$, and $(\partial_1 u_0)|_{\partial\Omega \cap U}$. We compute

$$\begin{aligned} \partial_1 A_{11} &= \partial_1 \gamma |\nabla u_0|^{p-2} \left(1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) \\ &\quad + \partial_1^2 u_0 \gamma \partial_1 u_0 |\nabla u_0|^{p-4} (p-2) \left(3 + (p-4) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) + a_{11}. \end{aligned} \quad (61)$$

For $j \neq 1$

$$\begin{aligned} \partial_1 A_{jj} &= \partial_1 \gamma |\nabla u_0|^{p-2} \left(1 + (p-2) \frac{(\partial_j u_0)^2}{|\nabla u_0|^2} \right) \\ &\quad + \partial_1^2 u_0 \gamma \partial_1 u_0 |\nabla u_0|^{p-4} (p-2) \left(1 + (p-4) \frac{(\partial_j u_0)^2}{|\nabla u_0|^2} \right) + a_{jj}. \end{aligned} \quad (62)$$

Also

$$\begin{aligned}\partial_1 A_{j1} &= \partial_1 \gamma |\nabla u_0|^{p-2} (p-2) \frac{\partial_j u_0 \partial_1 u_0}{|\nabla u_0|^2} \\ &\quad + \partial_1^2 u_0 \gamma \partial_j u_0 |\nabla u_0|^{p-4} (p-2) \left(1 + (p-4) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) + a_{j1}.\end{aligned}\quad (63)$$

Since $\nabla \cdot (\gamma |\nabla u_0|^{p-2} \nabla u_0) = 0$, at z we have that

$$\partial_1 [\gamma |\nabla u_0|^{p-2} \partial_1 u_0] = -\nabla' \cdot [\gamma |\nabla u_0|^{p-2} \nabla' u_0] = \partial_1 [\tilde{\gamma} |\nabla \tilde{u}_0|^{p-2} \partial_1 \tilde{u}_0], \quad (64)$$

by the previous step. Let $\xi_1 = \partial_1(\gamma - \tilde{\gamma})(z)$ and $\xi_2 = \partial_1^2(u_0 - \tilde{u}_0)(z)$. It follows that

$$\Theta_{11}\xi_1 + \Theta_{12}\xi_2 = 0, \quad (65)$$

where

$$\Theta_{11} = \partial_1 u_0 |\nabla u_0|^{p-2}, \quad (66)$$

$$\Theta_{12} = \gamma |\nabla u_0|^{p-2} \left(1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right). \quad (67)$$

Let $\xi_3 = \partial_1^2 \Phi^1(z)$. Taking $j = k = 1$ in (60), we obtain the equation

$$\Theta_{21}\xi_1 + \Theta_{22}\xi_2 + \Theta_{23}\xi_3 = 0, \quad (68)$$

where

$$\Theta_{21} = |\nabla u_0|^{p-2} \left(1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) \quad (69)$$

$$\Theta_{22} = \gamma \partial_1 u_0 |\nabla u_0|^{p-4} (p-2) \left(3 + (p-4) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) \quad (70)$$

$$\Theta_{23} = \gamma |\nabla u_0|^{p-2} \left(1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right). \quad (71)$$

Taking $k = j$ in (60), we obtain the equation

$$\Theta_{31}\xi_1 + \Theta_{32}\xi_2 + \Theta_{33}\xi_3 = -2\partial_1^2 \Phi^j \gamma |\nabla u_0|^{p-2} (p-2) \frac{\partial_j u_0 \partial_1 u_0}{|\nabla u_0|^2}, \quad (72)$$

where

$$\Theta_{31} = |\nabla u_0|^{p-2} \left(1 + (p-2) \frac{(\partial_j u_0)^2}{|\nabla u_0|^2} \right), \quad (73)$$

$$\Theta_{32} = \gamma \partial_1 u_0 |\nabla u_0|^{p-4} (p-2) \left(1 + (p-4) \frac{(\partial_j u_0)^2}{|\nabla u_0|^2} \right), \quad (74)$$

$$\Theta_{33} = -\gamma |\nabla u_0|^{p-2} \left(1 + (p-2) \frac{(\partial_j u_0)^2}{|\nabla u_0|^2} \right), \quad (75)$$

Under our assumptions, we are still free to rotate the coordinate axes, as long as the normal direction remains that of x_1 . We can therefore arrange that $\partial_j u_0(z) = 0$. In this case we then have the system

$$\begin{cases} \Theta_{11}\xi_1 + \Theta_{12}\xi_2 = 0, \\ \Theta_{21}\xi_1 + \Theta_{22}\xi_2 + \Theta_{23}\xi_3 = 0, \\ \Theta_{31}\xi_1 + \Theta_{32}\xi_2 + \Theta_{33}\xi_3 = 0. \end{cases} \quad (76)$$

Denoting $\lambda(z) = \gamma^2(z)|\nabla u_0|^{3p-8}(z)$, we compute the determinant

$$\begin{aligned} & \begin{vmatrix} \Theta_{11} & \Theta_{12} & 0 \\ \Theta_{21} & \Theta_{22} & \Theta_{23} \\ \Theta_{31} & \Theta_{32} & \Theta_{33} \end{vmatrix} \\ &= \lambda(z) \begin{vmatrix} \partial_1 u_0 & |\nabla u_0|^2 \left(1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) & 0 \\ 1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} & (p-2)\partial_1 u_0 \left(3 + (p-4) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) & 1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \\ 1 & (p-2)\partial_1 u_0 & -1 \end{vmatrix} \\ &= \lambda(z) \begin{vmatrix} \partial_1 u_0 & |\nabla u_0|^2 \left(1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) & 0 \\ 0 & (p-2)\partial_1 u_0 \left(3 + (p-4) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) & 1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \\ 2 & (p-2)\partial_1 u_0 & -1 \end{vmatrix} \\ &= \lambda(z) \left[2|\nabla u_0|^2 \left(1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right)^2 \right. \\ & \quad \left. - (p-2)(\partial_1 u_0)^2 \left(3 + (p-4) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) \right] \\ &= \lambda(z) \left[2|\nabla u_0|^2 + (p-2)(\partial_1 u_0)^2 + p(p-2) \frac{(\partial_1 u_0)^4}{|\nabla u_0|^2} \right] \neq 0, \end{aligned} \quad (77)$$

where the conclusion holds because, since $p > 1$, both $p-2 > -1$ and $p(p-2) > -1$. It follows that

$$\partial_1 \gamma(z) = \partial_1 \tilde{\gamma}(z), \quad \partial_1^2 u_0(z) = \partial_1^2 \tilde{u}_0(z), \quad \partial_1^2 \Phi^1(z) = 0. \quad (78)$$

Returning to (60), with $k = 1$, we are left with

$$A_{11} \partial_1^2 \Phi^j = 0, \quad (79)$$

which implies that

$$\partial_1^2 \Phi^j(z) = 0, \quad (80)$$

for all directions j that are orthogonal to the projection of ∇u_0 into the tangent plane. If we choose our coordinates so that the direction of x_l is the same as that of the just-mentioned projection, then $A_{l1}(z) \neq 0$, so (60), with $j = k = l$ gives

$$\partial_1^2 \Phi^l(z) = 0. \quad (81)$$

Therefore, combining (59) and (81), we have that

$$\partial_j \partial_k \Phi(z) = 0, \quad j, k = 1, \dots, n. \quad (82)$$

***m*-th order:**

For multiindices $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$, suppose that we know that

$$\partial^\alpha \gamma(z) = \partial^\alpha \tilde{\gamma}(z), \quad \alpha_1 = 0, 1, \dots, m-1, \quad (83)$$

$$\partial^\alpha u_0(z) = \partial^\alpha \tilde{u}_0(z), \quad \alpha_1 = 0, 1, \dots, m, \quad (84)$$

$$\partial^\alpha D\Phi(z) = \partial^\alpha I, \quad \alpha_1 = 0, 1, \dots, m-1. \quad (85)$$

We have that

$$(\partial_1^m \tilde{A})(z) = \left(\partial_1^m \frac{1}{|D\Phi|} (D\Phi)^T A D\Phi \right)(z). \quad (86)$$

Using our induction assumptions, we can rewrite this as

$$\partial_1^m \tilde{A}_{jk} = \partial_1^m A_{jk} + (A_{j1} \partial_1^{m+1} \Phi^k + A_{1k} \partial_1^{m+1} \Phi^j) - A_{jk} \partial_1^{m+1} \Phi^1. \quad (87)$$

Denoting by a_{jk} terms made up of quantities whose uniqueness follows from the induction hypotheses, that is, on the quantities in (83) and (83), we have that

$$\begin{aligned} \partial_1^m A_{jk} &= \partial_1^m \gamma(z) |\nabla u_0|^{p-2} \left(\delta_{jk} + (p-2) \frac{\partial_j u_0 \partial_k u_0}{|\nabla u_0|^2} \right) \\ &\quad + \partial_1^{m+1} u_0 (p-2) \gamma(z) |\nabla u_0|^{p-4} \\ &\quad \times \left(\delta_{jk} \partial_1 u_0 + (p-4) \frac{\partial_j u_0 \partial_k u_0}{|\nabla u_0|^2} \partial_1 u_0 + \delta_{1j} \partial_k u_0 + \delta_{1k} \partial_j u_0 \right) + a_{jk}. \end{aligned} \quad (88)$$

Since $\nabla \cdot (\gamma |\nabla u_0|^{p-2} \nabla u_0) = 0$, at z we have that

$$\begin{aligned} \partial_1^m [\gamma |\nabla u_0|^{p-2} \partial_1 u_0] \\ = -\partial_1^{m-1} \nabla' \cdot [\gamma |\nabla u_0|^{p-2} \nabla' u_0] = \partial_1^m [\tilde{\gamma} |\nabla \tilde{u}_0|^{p-2} \partial_1 \tilde{u}_0]. \end{aligned} \quad (89)$$

This can be rewritten as

$$\begin{aligned} \partial_1^m (\gamma - \tilde{\gamma}) \partial_1 u_0 |\nabla u_0|^{p-2} \\ + \partial_1^{m+1} (u_0 - \tilde{u}_0) \gamma |\nabla u_0|^{p-2} \left(1 + (p-2) \frac{(\partial_1 u_0)^2}{|\nabla u_0|^2} \right) = 0. \end{aligned} \quad (90)$$

If we set $\xi_1 = \partial_1^m (\gamma - \tilde{\gamma})(z)$, $\xi_2 = \partial_1^{m+1} (u_0 - \tilde{u}_0)$, $\xi_3 = \partial_1^{m+1} \Phi^1$, and if we choose a direction j that is orthogonal to the projection of $\nabla u_0(z)$ into the tangent space to $\partial\Omega$ at z , we obtain the same system

$$\begin{cases} \Theta_{11} \xi_1 + \Theta_{12} \xi_2 = 0, \\ \Theta_{21} \xi_1 + \Theta_{22} \xi_2 + \Theta_{23} \xi_3 = 0, \\ \Theta_{31} \xi_1 + \Theta_{32} \xi_2 + \Theta_{33} \xi_3 = 0. \end{cases} \quad (91)$$

Since with our assumptions

$$\begin{vmatrix} \Theta_{11} & \Theta_{12} & 0 \\ \Theta_{21} & \Theta_{22} & \Theta_{23} \\ \Theta_{31} & \Theta_{32} & \Theta_{33} \end{vmatrix} \neq 0, \quad (92)$$

it follows that

$$\partial_1^m \gamma(z) = \partial_1^m \gamma(z), \quad \partial_1^{m+1} u_0(z) = \partial_1^{m+1} u_0(z), \quad \partial_1^{m+1} \Phi^1(z) = 0. \quad (93)$$

Setting $k = 1$ in (87), we have

$$A_{11} \partial_1^{m+1} \Phi^j = 0, \quad (94)$$

which implies that

$$\partial_1^{m+1} \Phi^j(z) = 0, \quad (95)$$

for all directions j that are orthogonal to the projection of ∇u_0 into the tangent plane. If we choose our coordinates so that the direction of x_l is the same as that of the just-mentioned projection, then $A_{l1}(z) \neq 0$, so (87), with $j = k = l$ gives

$$\partial_1^{m+1} \Phi^l(z) = 0. \quad (96)$$

Therefore

$$\partial^\alpha D\Phi(z) = \partial^\alpha I, \quad \alpha_1 = 0, 1, \dots, m. \quad (97)$$

This completes the induction step.

Competing interest. The authors have no competing interests to declare.

Financial support. C.C. was supported by NSTC grants 112-2115-M-A49-002 and 113-2115-M-A49-018-MY3.

References

- [1] G. S. Alberti, G. Bal and M. Di Cristo, ‘Critical points for elliptic equations with prescribed boundary conditions’, *Arch. Ration. Mech. Anal.*, **226** (2017), 117–141.
- [2] G. Alessandrini and M. Sigalotti, ‘Geometric properties of solutions to the anisotropic p-Laplace equation in dimension two’, *Ann. Fenn. Math.*, **26**(1) (2001), 249–266.
- [3] T. Brander, ‘Calderón problem for the p-Laplacian: First order derivative of conductivity on the boundary’, *Proc. Amer. Math. Soc.*, **144**(1) (2016), 177–189.
- [4] T. Brander, B. Harrach, M. Kar and M. Salo, ‘Monotonicity and enclosure methods for the p-Laplace equation’, *SIAM J. Appl. Math.*, **78**(2) (2018), 742–758.
- [5] T. Brander, J. Ilmavirta and M. Kar, ‘Superconductive and insulating inclusions for linear and non-linear conductivity equations’, *Inverse Probl. Imaging*, **12**(1) (2018), 91–123.
- [6] T. Brander, M. Kar and M. Salo, ‘Enclosure method for the p-Laplace equation’, *Inverse Probl.*, **31**(4) (2015), 045001.
- [7] A. P. Calderón, ‘On an inverse boundary value problem’, in *Seminar on Numerical Analysis and Its Applications to Continuum Physics* (Rio de Janeiro, 1980), 65–73 (Soc. Brasil. Mat., Rio de Janeiro, 1980).
- [8] C. I. Cârstea, ‘On an inverse boundary value problem for a nonlinear time harmonic Maxwell system’, *J. Inverse Ill-posed Probl.*, **30**(3) (2022), 395–408.
- [9] C. I. Cârstea and A. Feizmohammadi, ‘A density property for tensor products of gradients of harmonic functions and applications’, Preprint, 2020, [arXiv:2009.11217](https://arxiv.org/abs/2009.11217).
- [10] C. I. Cârstea and A. Feizmohammadi, ‘An inverse boundary value problem for certain anisotropic quasilinear elliptic equations’, *J. Differ. Equ.*, **284** (2021), 318–349.
- [11] C. I. Cârstea, A. Feizmohammadi, Y. Kian, K. Krupchyk and G. Uhlmann, ‘The Calderón inverse problem for isotropic quasilinear conductivities’, *Adv. Math.*, **391** (2021), 107956.
- [12] C. I. Cârstea, T. Ghosh and G. Nakamura, ‘An inverse boundary value problem for the inhomogeneous porous medium equation’, Preprint, 2021, [arXiv:2105.01368](https://arxiv.org/abs/2105.01368).
- [13] C. I. Cârstea, T. Ghosh and G. Uhlmann, ‘An inverse problem for the porous medium equation with partial data and a possibly singular absorption term’, *SIAM J. Math. Anal.*, **55**(1) (2023), 162–185.
- [14] C. I. Cârstea and M. Kar, ‘Recovery of coefficients for a weighted p-Laplacian perturbed by a linear second order term’, *Inverse Probl.*, **37**(1) (2020), 015013.

- [15] C. I. Cârstea, G. Nakamura and M. Vashisth, 'Reconstruction for the coefficients of a quasilinear elliptic partial differential equation', *Appl. Math. Lett.*, **98** (2019), 371–377.
- [16] H. Egger, J.-F. Pietschmann and M. Schlottbom, 'Simultaneous identification of diffusion and absorption coefficients in a quasilinear elliptic problem', *Inverse Probl.*, **30**(3) (2014), 035009.
- [17] A. Feizmohammadi and L. Oksanen, 'An inverse problem for a semi-linear elliptic equation in Riemannian geometries', *J. Differ. Equ.*, **269**(6) (2020), 4683–4719.
- [18] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, *Classics in Mathematics*, 224 (Springer, Berlin, 1998).
- [19] C.-Y. Guo, M. Kar and M. Salo, 'Inverse problems for p-Laplace type equations under monotonicity assumptions', *Rend. Istit. Mat. Univ. Trieste*, **48** (2016), 79–99.
- [20] D. Hervas and Z. Sun, 'An inverse boundary value problem for quasilinear elliptic equations', *Commun. Partial Differ. Equ.*, **27**(11–12) (2002), 2449–2490.
- [21] V. Isakov, 'On uniqueness in inverse problems for semilinear parabolic equations', *Arch. Ration. Mech. Anal.*, **124**(1) (1993), 1–12.
- [22] V. Isakov, 'Uniqueness of recovery of some quasilinear partial differential equations', *Commun. Partial Differential Equations* **26**(11–12) (2001), 1947–1973.
- [23] V. Isakov and A. I. Nachman, 'Global uniqueness for a two-dimensional semilinear elliptic inverse problem', *Trans. Amer. Math. Soc.* **347**(9) (1995), 3375–3390.
- [24] V. Isakov and J. Sylvester, 'Global uniqueness for a semilinear elliptic inverse problem', *Commun. Pure Appl. Math.* **47**(10) (1994), 1403–1410.
- [25] H. Kang and G. Nakamura, 'Identification of nonlinearity in a conductivity equation via the Dirichlet-to-Neumann map', *Inverse Problems* **18**(4) (2002), 1079.
- [26] H. Kang and K. Yun, 'Boundary determination of conductivities and Riemannian metrics via local Dirichlet-to-Neumann operator', *SIAM J. Math. Anal.* **34**(3) (2002), 719–735.
- [27] M. Kar and J.-N. Wang, 'Size estimates for the weighted p-Laplace equation with one measurement', *Discrete Contin. Dyn. Syst. B* **22**(11) (2017).
- [28] K. Krupchyk and G. Uhlmann, 'Partial data inverse problems for semilinear elliptic equations with gradient nonlinearities', *Math. Res. Lett.* **27**(6) (2020).
- [29] K. Krupchyk and G. Uhlmann, 'A remark on partial data inverse problems for semilinear elliptic equations', *Proc. Amer. Math. Soc.* **148**(2) (2020), 681–685.
- [30] M. Lassas, T. Liimatainen, Y.-H. Lin, and M. Salo, 'Partial data inverse problems and simultaneous recovery of boundary and coefficients for semilinear elliptic equations', *Rev. Mat. Iberoam.* **37**(4) (2020), 1553–1580.
- [31] M. Lassas, T. Liimatainen, Y.-H. Lin, and M. Salo, 'Inverse problems for elliptic equations with power type nonlinearities', *J. Math. Pures Appl.* **145** (2021), 44–82.
- [32] J. M. Lee and G. Uhlmann, 'Determining anisotropic real-analytic conductivities by boundary measurements', *Commun. Pure Appl. Math.* **42**(8) (1989), 1097–1112.
- [33] G. M. Lieberman, 'Boundary regularity for solutions of degenerate elliptic equations', *Nonlinear Anal.* **12**(11) (1988), 1203–1219.
- [34] C. Munoz and G. Uhlmann, 'The Calderón problem for quasilinear elliptic equations', *Ann. Inst. H. Poincaré C Anal. Non Linéaire* (2020).
- [35] A. Nachman, 'Global uniqueness for a two-dimensional inverse boundary value problem', *Ann. of Math.* **143**(1) (1996), 71–96.
- [36] M. Salo and X. Zhong, 'An inverse problem for the p-Laplacian: boundary determination', *SIAM J. Math. Anal.* **44**(4) (2012), 2474–2495.
- [37] R. Shankar, 'Recovering a quasilinear conductivity from boundary measurements', *Inverse Problems* **37**(1) (2020), 015014.
- [38] Z. Sun, 'On a quasilinear inverse boundary value problem', *Math. Z.* **221**(1) (1996), 293–305.
- [39] Z. Sun, 'Anisotropic inverse problems for quasilinear elliptic equations', *J. Phys. Conf. Ser.* **12** (2005), 015.
- [40] Z. Sun, 'An inverse boundary-value problem for semilinear elliptic equations', *Electron. J. Differential Equations* (2010), Paper No.
- [41] Z. Sun and G. Uhlmann, 'Inverse problems in quasilinear anisotropic media', *Amer. J. Math.* **119**(4) (1997), 771–797.
- [42] J. Sylvester and G. Uhlmann, 'A global uniqueness theorem for an inverse boundary value problem', *Ann. of Math.* **125**(1) (1987), 153–169.