

A TERMINATING INTUITIONISTIC CALCULUS

GIULIO FELLIN¹ AND SARA NEGRI

Abstract. A terminating sequent calculus for intuitionistic propositional logic is obtained by modifying the $R\supset$ rule of the labelled sequent calculus **G3I**. This is done by adding a variant of the principle of *a fortiori* in the left-hand side of the premiss of the rule. In the resulting calculus, called **G3I_f**, derivability of any given sequent is directly decidable by root-first proof search, without any extra device such as loop-checking. In the negative case, the failed proof search gives a finite countermodel to the sequent on a reflexive, transitive, and Noetherian Kripke frame. As a byproduct, a direct proof of faithfulness of the embedding of intuitionistic logic into Grzegorczyk logic is obtained.

§1. Introduction. In his doctoral thesis [8, 9], Gentzen introduced sequent calculi for classical and intuitionistic logic. In particular, he solved the decision problem for intuitionistic propositional logic (**Int**) with a calculus that he called **LI**.¹ However, Gentzen’s original calculus lacked some desirable properties, such as invertibility of its rules, that would avoid the need for backtracking. Ever since then, many other approaches were proposed; we refer to [4] for an extended survey.

The labelled calculus **G3I** by Dyckhoff and Negri [5, 14, 17] reported in Table 1 solves the problem of backtracking but doesn’t yet have the property of termination, see, for instance, the example of Peirce’s Law in Section 3.3. In order to solve this problem, Negri [15, 16] showed how to add a loop-checking mechanism to ensure termination. However, it is desirable to avoid loop-checking since its effect on complexity is not clear.

Corsi [2, 3] presented a calculus for **Int** which fulfils the termination property. The key to get termination is the addition of the following rule:

$$\frac{\Gamma \rightarrow \Delta, B}{\Gamma \rightarrow \Delta, A \supset B} \text{a fortiori}.$$

This rule is logically equivalent to the formula $B \supset (A \supset B)$, which is called the principle of *a fortiori*.

Received October 5, 2022.

2020 *Mathematics Subject Classification*. Primary 03B20, 03F07, Secondary 03B45, 03F45.

Key words and phrases. intuitionistic logic, proof theory, sequent calculus, Grzegorczyk logic.

¹To clarify a point raised by an anonymous referee, we note that Gentzen designated his intuitionistic calculus by ‘NI’. His handwriting for capital ‘I’ was in the old Sütterlin handwriting that has been rendered as ‘J’ in the printing, which contemporary readers of his article would have understood. These practices have been communicated to us by Jan von Plato. For the Sütterlin writing of capital ‘I’, the one that in today’s eyes resembles ‘J’, see [19, p. 87].

© The Author(s), 2023. Published by Cambridge University Press on behalf of The Association for Symbolic Logic. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

0022-4812/25/9001-0011

DOI:10.1017/jsl.2023.88

In this paper, we consider the labelled calculus **G3I** instead, and show that a way to reach termination consists in modifying its rule $R\supset$ as follows:

$$\frac{x \leq y, y: B \supset (A \supset B), y: A, \Gamma \rightarrow \Delta, y: B}{\Gamma \rightarrow \Delta, x: A \supset B} R\supset_t \quad (y \text{ fresh}).$$

Although the idea comes from a similar terminating procedure [6] for the calculus **G3Grz** for the provability logic **Grz**, into which **Int** is embeddable as detailed in Section 4, we notice that what we do is actually incorporating *a fortiori* into $R\supset$.

§2. Structural properties. Consider the sequent calculi **G3I** and **G3I_t** as presented in Tables 1 and 2, respectively.

Initial sequent

$$x \leq y, x: P, \Gamma \rightarrow \Delta, y: P$$

Logical rules

$$\frac{x: A, x: B, \Gamma \rightarrow \Delta}{x: A \wedge B, \Gamma \rightarrow \Delta} L\wedge$$

$$\frac{\Gamma \rightarrow \Delta, x: A \quad \Gamma \rightarrow \Delta, x: B}{\Gamma \rightarrow \Delta, x: A \wedge B} R\wedge$$

$$\frac{x: A, \Gamma \rightarrow \Delta \quad x: B, \Gamma \rightarrow \Delta}{x: A \vee B, \Gamma \rightarrow \Delta} L\vee$$

$$\frac{\Gamma \rightarrow \Delta, x: A, x: B}{\Gamma \rightarrow \Delta, x: A \vee B} R\vee$$

$$\frac{x \leq y, x: A \supset B, \Gamma \rightarrow \Delta, y: A \quad x \leq y, x: A \supset B, y: B, \Gamma \rightarrow \Delta}{x \leq y, x: A \supset B, \Gamma \rightarrow \Delta} L\supset$$

$$\frac{}{x: \perp, \Gamma \rightarrow \Delta} L\perp$$

$$\frac{x \leq y, y: A, \Gamma \rightarrow \Delta, y: B}{\Gamma \rightarrow \Delta, x: A \supset B} R\supset$$

Mathematical rules

$$\frac{x \leq x, \Gamma \rightarrow \Delta}{\Gamma \rightarrow \Delta} \text{Ref}_{\leq}$$

$$\frac{x \leq z, x \leq y, y \leq z, \Gamma \rightarrow \Delta}{x \leq y, y \leq z, \Gamma \rightarrow \Delta} \text{Trans}_{\leq}$$

TABLE 1. The sequent calculus **G3I**. Rule $R\supset$ has the condition that y is fresh.

Initial sequent

As in **G3I**

Logical rules

$L\wedge, R\wedge, L, R, L\supset, L\perp$ as in **G3I**

$$\frac{x \leq y, y: B \supset (A \supset B), y: A, \Gamma \rightarrow \Delta, y: B}{\Gamma \rightarrow \Delta, x: A \supset B} R\supset_t \quad (y \text{ fresh})$$

Mathematical rules

As in **G3I**

TABLE 2. The sequent calculus **G3I_t**.

THEOREM 2.1. **G3I** and **G3I_t** are equivalent in the sense that

$$\mathbf{G3I} \vdash \Gamma \rightarrow \Delta \quad \text{if and only if} \quad \mathbf{G3I}_t \vdash \Gamma \rightarrow \Delta.$$

PROOF. Suppose that $\mathbf{G3I} \vdash \Gamma \rightarrow \Delta$. We transform the given derivation into one in $\mathbf{G3I}_t$ by using height-preserving weakening to add whenever needed the extra formula of the form $y: B \supset (A \supset B)$ in the premiss of $R\supset$. So $\mathbf{G3I}_t \vdash \Gamma \rightarrow \Delta$.

Conversely, if $\mathbf{G3I}_t \vdash \Gamma \rightarrow \Delta$, consider the steps of $R\supset_t$:

$$\frac{x \leq y, y: B \supset (A \supset B), y: A, \Gamma' \rightarrow \Delta', y: B}{\Gamma' \rightarrow \Delta', x: A \supset B} R\supset_t$$

$$\vdots$$

$$\Gamma \rightarrow \Delta.$$

By a Cut with the extra (derivable) sequent $\rightarrow y: B \supset (A \supset B)$, we turn it into premisses of $R\supset$ with the same conclusions:

$$\frac{\rightarrow y: B \supset (A \supset B) \quad x \leq y, y: B \supset (A \supset B), y: A, \Gamma' \rightarrow \Delta', y: B}{x \leq y, y: A, \Gamma' \rightarrow \Delta', y: B} \text{Cut}$$

$$\frac{x \leq y, y: A, \Gamma' \rightarrow \Delta', y: B}{\Gamma' \rightarrow \Delta', x: A \supset B} R\supset$$

$$\vdots$$

$$\Gamma \rightarrow \Delta.$$

We conclude by admissibility of Cut in **G3I**. ⊢

THEOREM 2.2. All the structural properties hold for **G3I_t**. In particular,

- (i) All the sequents of the following form are derivable in **G3I_t**:
 - (a) $x \leq y, x: A, \Gamma \rightarrow \Delta, y: A$,
 - (b) $x: A, \Gamma \rightarrow \Delta, x: A$.
- (ii) If $\mathbf{G3I}_t \vdash \Gamma \rightarrow \Delta$, then $\mathbf{G3I}_t \vdash \Gamma(x/y) \rightarrow \Delta(x/y)$ with the same derivation height.
- (iii) The rules of weakening,

$$\frac{\Gamma \rightarrow \Delta}{x: A, \Gamma \rightarrow \Delta} LW \quad \frac{\Gamma \rightarrow \Delta}{\Gamma \rightarrow \Delta, x: A} RW \quad \frac{\Gamma \rightarrow \Delta}{x \leq y, \Gamma \rightarrow \Delta} LW_{\leq}$$

are height-preserving admissible in **G3I_t**.

- (iv) All rules of **G3I_t** are height-preserving invertible.
- (v) The rules of contraction,

$$\frac{x: A, x: A, \Gamma \rightarrow \Delta}{x: A, \Gamma \rightarrow \Delta} LC \quad \frac{\Gamma \rightarrow \Delta, x: A, x: A}{\Gamma \rightarrow \Delta, x: A} RC$$

$$\frac{x \leq y, x \leq y, \Gamma \rightarrow \Delta}{x \leq y, \Gamma \rightarrow \Delta} LC_{\leq}$$

are height-preserving admissible in **G3I_t**.

PROOF. The proofs are similar to those of [17, Lemmas 12.25–12.29 and Theorems 12.27–12.29]. ⊢

We will see later that also the rule of cut,

$$\frac{\Gamma \rightarrow \Delta, x : A \quad x : A, \Gamma' \rightarrow \Delta'}{\Gamma, \Gamma' \rightarrow \Delta, \Delta'} \text{Cut}$$

is admissible in $\mathbf{G3I}_t$ (Corollary 3.8). However, the proof that we are going to give is not syntactical. On the other hand, this formulation of the calculus permits a completeness proof (Theorem 3.5) that yields at the same time a semantic proof of admissibility of Cut, the finite model property and a constructive decision procedure.

REMARK 2.3. As a consequence of admissibility of LW, rule $R\supset$ of $\mathbf{G3I}$ is admissible in $\mathbf{G3I}_t$.

We now prove a few lemmata that will be useful later.

LEMMA 2.4. *The rule*

$$\frac{x \leq y, \Gamma \rightarrow \Delta, x : A}{x \leq y, \Gamma \rightarrow \Delta, y : A}$$

is admissible in $\mathbf{G3I}_t$.

PROOF. We prove it by induction on the height of the derivation of the premiss.

$n = 0$: The only nontrivial case is the one in which the premiss is an initial sequent and $x : A$ is principal. In this case, we can write the sequent as

$$x \leq y, w \leq x, w : A, \Gamma' \rightarrow \Delta, x : A,$$

where $\Gamma \equiv w \leq x, w : A, \Gamma'$. Observe that the sequent

$$w \leq y, x \leq y, w \leq x, w : A, \Gamma' \rightarrow \Delta, y : A$$

is initial. By the rule $\text{Trans}_{\leq}$, we get a derivation of

$$x \leq y, w \leq x, w : A, \Gamma' \rightarrow \Delta, y : A,$$

which is just $x \leq y, \Gamma \rightarrow \Delta, y : A$, as wanted.

$n > 0$: The only nontrivial cases are those in which the last rule applied is a right rule and $x : A$ is principal. If the last rule applied is $R\wedge$ and $A \equiv B \wedge C$, then we have

$$\frac{x \leq y, \Gamma \rightarrow \Delta, x : B \quad x \leq y, \Gamma \rightarrow \Delta, x : C}{x \leq y, \Gamma \rightarrow \Delta, x : B \wedge C} R\wedge.$$

We can apply the induction hypothesis to the premisses and get

$$\begin{aligned} x &\leq y, \Gamma \rightarrow \Delta, y : B, \\ x &\leq y, \Gamma \rightarrow \Delta, y : C. \end{aligned}$$

We conclude by an application of $R\wedge$. If the last rule applied is $R\vee$ and $A \equiv B \vee C$, then we have

$$\frac{x \leq y, \Gamma \rightarrow \Delta, x : B, x : C}{x \leq y, \Gamma \rightarrow \Delta, x : B \vee C} R\vee.$$

We can apply the induction hypothesis to the premiss and get

$$x \leq y, \Gamma \rightarrow \Delta, y : B, y : C.$$

We conclude by an application of $R\supset$. If the last rule applied is $R\supset_t$ and $A \equiv B \supset C$, then we have

$$\frac{x \leq z, x \leq y, z : C \supset (B \supset C), z : B, \Gamma \rightarrow \Delta, z : C}{x \leq y, \Gamma \rightarrow \Delta, x : B \supset C} R\supset_t.$$

We can apply hp-weakening to the premiss and get

$$y \leq z, x \leq z, x \leq y, z : C \supset (B \supset C), z : B, \Gamma \rightarrow \Delta, z : C,$$

which, by an application of transitivity leads to

$$y \leq z, x \leq y, z : C \supset (B \supset C), z : B, \Gamma \rightarrow \Delta, z : C.$$

We conclude with an application of $R\supset_t$. ⊢

LEMMA 2.5. *The rule*

$$\frac{x \leq y, x : A, y : A, \Gamma \rightarrow \Delta}{x \leq y, x : A, \Gamma \rightarrow \Delta}$$

is admissible in $\mathbf{G3I}_t$.

PROOF. We prove it by induction on the height of the derivation of the premiss.

$n = 0$: The only nontrivial case is the one in which the premiss is an initial sequent and $y : A$ is principal. In this case, we can write the sequent as

$$x \leq y, y \leq z, x : A, y : A, \Gamma' \rightarrow \Delta', z : A,$$

where $\Gamma \equiv y \leq z, \Gamma'$ and $\Delta \equiv \Delta', z : A$. Observe that the sequent

$$x \leq y, y \leq z, x \leq z, x : A, \Gamma' \rightarrow \Delta', z : A$$

is initial. By transitivity, we get a derivation of

$$x \leq y, y \leq z, x : A, \Gamma' \rightarrow \Delta', z : A,$$

which is just $x \leq y, x : A, \Gamma \rightarrow \Delta$, as wanted.

$n > 0$: The only nontrivial cases are those in which the last rule applied is a left rule and $y : A$ is principal. If the last rule applied is $L\wedge$ and $A \equiv B \wedge C$, then we have

$$\frac{x \leq y, x : B \wedge C, y : B, y : C, \Gamma \rightarrow \Delta}{x \leq y, x : B \wedge C, y : B \wedge C, \Gamma \rightarrow \Delta} L\wedge.$$

Then, by hp-invertibility of $L\wedge$, we get

$$x \leq y, x : B, x : C, y : B, y : C, \Gamma \rightarrow \Delta,$$

to which the induction hypothesis can be applied:

$$x \leq y, x : B, x : C, \Gamma \rightarrow \Delta.$$

We conclude by an application of $L\wedge$. If the last rule applied is $L\vee$ and $A \equiv B \vee C$, then we have

$$\frac{x \leq y, x : B \vee C, y : B, \Gamma \rightarrow \Delta \quad x \leq y, x : B \vee C, y : C, \Gamma \rightarrow \Delta}{x \leq y, x : B \vee C, y : B \vee C, \Gamma \rightarrow \Delta} L\vee.$$

Then, by hp-invertibility of $L\vee$, we get

$$\begin{aligned} x &\leq y, x : B, y : B, \Gamma \rightarrow \Delta, \\ x &\leq y, x : C, y : C, \Gamma \rightarrow \Delta, \end{aligned}$$

to which the induction hypothesis can be applied:

$$\begin{aligned} x &\leq y, x : B, \Gamma \rightarrow \Delta, \\ x &\leq y, x : C, \Gamma \rightarrow \Delta. \end{aligned}$$

We conclude by an application of $L\supset$. If the last rule applied is $L\supset$ and $A \equiv B \supset C$, then we have

$$\frac{x \leq y, x : B \supset C, y : B \supset C, y \leq z, \Gamma' \rightarrow \Delta, z : B \quad x \leq y, x : B \supset C, y : B \supset C, z : C, y \leq z, \Gamma' \rightarrow \Delta}{x \leq y, x : B \supset C, y : B \supset C, y \leq z, \Gamma' \rightarrow \Delta} L\supset,$$

where $\Gamma \equiv y \leq z, \Gamma'$. Then we can apply the induction hypothesis to the premisses:

$$\begin{aligned} x &\leq y, x : B \supset C, y \leq z, \Gamma' \rightarrow \Delta, z : B, \\ x &\leq y, x : B \supset C, z : C, y \leq z, \Gamma' \rightarrow \Delta. \end{aligned}$$

By hp-weakening, these lead to

$$\begin{aligned} x &\leq z, x \leq y, x : B \supset C, y \leq z, \Gamma' \rightarrow \Delta, z : B, \\ x &\leq z, x \leq y, x : B \supset C, z : C, y \leq z, \Gamma' \rightarrow \Delta. \end{aligned}$$

Now we can apply $L\supset$ in order to get

$$x \leq z, x \leq y, x : B \supset C, y \leq z, \Gamma' \rightarrow \Delta.$$

We conclude by an application of transitivity. $\dashv$

LEMMA 2.6. *The rule*

$$\frac{x \leq y, x : B \supset (A \supset B), x : A, \Gamma \rightarrow \Delta, y : B, y : A \supset B}{x \leq y, x : B \supset (A \supset B), x : A, \Gamma \rightarrow \Delta, y : B}$$

is admissible.

PROOF. The direction from conclusion to premiss is just an instance of weakening. For the other direction, we apply invertibility of $R\supset_i$ (we notice that the inverse rule does not have the condition on the eigenvariable, but it can be done on an arbitrary label) to get

$$\frac{x \leq y, x \leq y, x : B \supset (A \supset B), y : B \supset (A \supset B), y : A, y : A, \Gamma \rightarrow \Delta, y : B, y : B}{x \leq y, x : B \supset (A \supset B), y : B \supset (A \supset B), x : A, y : A, \Gamma \rightarrow \Delta, y : B} LC, RC$$

Lemma 2.5, twice

$$x \leq y, x : B \supset (A \supset B), x : A, \Gamma \rightarrow \Delta, y : B.$$

$\dashv$

§3. Soundness and completeness.

3.1. Semantics. A *Kripke model* [11] (X, R, val) is a set X together with an *accessibility relation* R , i.e., a binary relation between elements of X , and a *valuation* val , i.e., a function assigning one of the truth values 0 or 1 to an element x of X and an atomic formula P . The usual notation for $\text{val}(x, P) = 1$ is $x \Vdash P$. In Kripke models for intuitionistic logic, the accessibility relation is a preorder, i.e., it is *reflexive*

$$\forall x (xRx)$$

and *transitive*

$$\forall x \forall y \forall z (zRy \ \& \ yRx \Rightarrow zRx),$$

and therefore it is denoted by the usual symbol $\leq$ for a preorder. For convenience, we assume to have equality $=$ and a binary relation $<$ on X which is transitive and *irreflexive*, i.e.,

$$\forall x (x \not< x),$$

and we define $\leq$ as its *reflexive closure*:

$$x \leq y \iff (x < y \text{ or } x = y).$$

As usual, we denote by $\geq$ the inverse relation of $\leq$; i.e.,

$$x \geq y \iff y \leq x.$$

The inductive definition of truth of a proposition in **Int** in terms of Kripke semantics is:

$$x \not\Vdash \perp.$$

$$x \Vdash A \wedge B \text{ if and only if } x \Vdash A \text{ and } x \Vdash B.$$

$$x \Vdash A \vee B \text{ if and only if } x \Vdash A \text{ or } x \Vdash B.$$

$$x \Vdash A \supset B \text{ if and only if } y \Vdash A \Rightarrow y \Vdash B \text{ for all } y \text{ such that } x \leq y.$$

Let $x \in X$. We say that $\leq$ *satisfies the semantic a fortiori property for x* if

$$\forall y \geq x (y \Vdash B \supset (A \supset B) \ \& \ y \Vdash A \Rightarrow y \Vdash B). \quad (\text{SAF}_x)$$

Let R be a relation on X . An *infinite R -sequence* is a sequence $(x_i)_{i \in \mathbb{N}}$ of elements of X such that $x_i R x_{i+1}$ for all $i \in \mathbb{N}$. An infinite R -sequence $(x_i)_{i \in \mathbb{N}}$ is *convergent* if there is $i \in \mathbb{N}$ such that $x_j = x_i$ for all $j > i$. We say that R is *Noetherian*—for short, R satisfies *Noeth*—if every infinite R -sequence converges.

LEMMA 3.1. *Let $x \in X$. If $\leq$ is Noetherian and satisfies SAF_x , then*

$$\forall y > x (y \Vdash B \supset (A \supset B)).$$

PROOF. Notice that the relation $<$ is transitive, irreflexive, and Noetherian. Therefore it follows that its inverse $>$ satisfies the Gödel–Löb Induction (see [7, Proposition 4.2 and Theorem 4.3]), that is,

$$\forall x (\forall y > x (\forall z > y Ez \Rightarrow Ey) \Rightarrow \forall y > x Ey) \quad (\text{GL-Ind})$$

for any given predicate $E(x)$ on X . Therefore, if we let $E(x) \equiv x \Vdash B \supset (A \supset B)$, it suffices to show that

$$\forall y > x (\forall z > y (z \Vdash B \supset (A \supset B)) \Rightarrow y \Vdash B \supset (A \supset B)). \quad (1)$$

So let $y > x$ such that

$$\forall z > y (z \Vdash B \supset (A \supset B)). \quad (2)$$

We claim that $y \Vdash B \supset (A \supset B)$, i.e.,

$$\forall z \geq y (z \Vdash B \Rightarrow z \Vdash A \supset B). \quad (3)$$

So let $z \geq y$ such that $z \Vdash B$. We have to prove $z \Vdash A \supset B$, i.e.,

$$\forall w \geq z (w \Vdash A \Rightarrow w \Vdash B). \quad (4)$$

So let $w \geq z$ such that $w \Vdash A$. The claim is $w \Vdash B$.

- If $w = z$, then we already know that $z \Vdash B$.
- If $w > z$, then by transitivity $w > y$ and by (2) we get $w \Vdash B \supset (A \supset B)$. Since $w \Vdash A$ and by transitivity $w \geq x$, we can apply SAF_x and derive $w \Vdash B$.

Now unroll the proof to get claims (4), (3), and (1), and thus the main claim. $\dashv$

LEMMA 3.2. *Fix $x \in X$. If $\leq$ is Noetherian and satisfies SAF_x , then $x \Vdash B \supset (A \supset B)$.*

PROOF. The claim is equivalent to

$$\forall y \geq x (y \Vdash B \Rightarrow y \Vdash A \supset B). \quad (5)$$

Fix $y \geq x$ such that $y \Vdash B$. We claim that $y \Vdash A \supset B$, i.e.,

$$\forall z \geq y (z \Vdash A \Rightarrow z \Vdash B). \quad (6)$$

Fix $z \geq y$ such that $z \Vdash A$. We need to prove that $z \Vdash B$.

- If $z = y$, then we already know that $y \Vdash B$.
- If $z > y$, then by transitivity $z > x$ and by Lemma 3.1 we get $z \Vdash B \supset (A \supset B)$. Since $z \Vdash A$ and by transitivity $z \geq x$, we can apply SAF_x and derive $z \Vdash B$.

Now unroll the proof to get claims (6) and (5), and thus the main claim. $\dashv$

LEMMA 3.3 (Semantic Lemma). *Fix $x \in X$. If $\leq$ is Noetherian, then the following are equivalent:*

- (i) SAF_x .
- (ii) $\forall y \geq x (y \Vdash A \Rightarrow y \Vdash B)$.

PROOF. (ii) $\Rightarrow$ (i): *A fortiori*.

(i) $\Rightarrow$ (ii): Fix $y \geq x$ such that $y \Vdash A$. We claim that $y \Vdash B$.

- If $y = x$, then by Lemma 3.2 we get that $x \Vdash B \supset (A \supset B)$.
- If $y > x$, then by Lemma 3.1 we get that $y \Vdash B \supset (A \supset B)$.

In either case we have $y \Vdash B \supset (A \supset B)$ and $y \Vdash A$, thus we can apply SAF_x and get $y \Vdash B$. $\dashv$

3.2. Proof search. Consider the proof search procedure as defined in [6]. We have the analogous of 5.3–6:

THEOREM 3.4 (Soundness). *If $\mathbf{G3I}_t \vdash \Gamma \rightarrow \Delta$, then $\Gamma \rightarrow \Delta$ is valid in every reflexive transitive and Noetherian frame.*

PROOF. If $\mathbf{G3I}_t \vdash \Gamma \rightarrow \Delta$, then $\mathbf{G3I} \vdash \Gamma \rightarrow \Delta$ and therefore $\Gamma \rightarrow \Delta$ is valid in every reflexive transitive frame, a fortiori in every Noetherian one. $\dashv$

THEOREM 3.5. *Let $\Gamma \rightarrow \Delta$ be a sequent in the language of $\mathbf{G3I}_t$. Then it is decidable whether it is derivable in $\mathbf{G3I}_t$. If it is not derivable, the failed proof search gives a finite countermodel to the sequent on a reflexive, transitive, and Noetherian frame.*

PROOF. We adapt the proof of [6, Theorem 5.4], which in turn is an adaptation to labelled sequents of the method of reduction trees detailed for Gentzen's LK by Takeuti [20, Chapter 1, Paragraph 8].

For an arbitrary sequent $\Gamma \rightarrow \Delta$ in the language of $\mathbf{G3I}_t$ we apply, whenever possible, root-first the rules of $\mathbf{G3I}_t$, in a given order. The procedure will construct either a derivation in $\mathbf{G3I}_t$ or a countermodel.

1. Construction of the reduction tree: The reduction tree is defined inductively in stages as follows: Stage 0 has $\Gamma \rightarrow \Delta$ at the root of the tree. For each branch, stage $n > 0$ has two cases:

Case I: If the top-sequent is either an initial sequent or has some $x : A$, not necessarily atomic, on both left and right, or is a conclusion of $L\perp$, the construction of the branch ends.

Case II: Otherwise we continue the construction of the branch by writing, above its top-sequent, other sequents that are obtained by applying root-first the rules of $\mathbf{G3I}_t$ (except $L\perp$) whenever possible, in a given order and under suitable conditions.

There are eight different stages: one for each logical rule, $\text{Ref}_{\leq}$ and Trans . At stage $8 + 1$ we repeat stage 1, at stage $8 + 2$ we repeat stage 2, and so on until an initial sequent, or a conclusion of $L\perp$, or a *saturated branch* (defined below) is found. In applying root-first the rules, we also copy their principal formulas in the premisses. All such copied formulas, except the principal formula of $L\supset$, need not be analysed again and are thus marked as overlined. For instance:

$$\frac{\overline{y : A \wedge B}, y : A, y : B, \Gamma \rightarrow \Delta}{y : A \wedge B, \Gamma \rightarrow \Delta} L\wedge$$

These marked formulas are only auxiliary, and will thus be removed at the end of the procedure to get the reduction tree.

The stages for the rules other than $R\supset_t$ are similar to those in [17, Theorem 11.28].

For formulas of the form $y : A \supset B$ in the succedent, we apply rule $R\supset_t$. However, if the sequent contains $x \leq y, x : B \supset (A \supset B), y : A$ in the antecedent and $y : B$ in the succedent, we remove it. This is justified by Lemma 2.6.

Finally, we consider the cases of the frame rules $\text{Ref}_{\leq}$ and Trans . As detailed in [5, 6], it is enough to instantiate $\text{Ref}_{\leq}$ only on terms in the top-sequent.

Observe also that, because of height-preserving admissibility of contraction, once a rule has been considered, it need not be instantiated again on the same principal formulas (for $L\supset$ such principal formulas are pairs of the form $x \leq y, x : A \supset B$ and

it need not be applied whenever its application produces a duplication of labelled formulas or relational atoms.

To show that the procedure terminates, it is enough to show that every branch in the reduction tree for a sequent $\Gamma \rightarrow \Delta$ is finite. Every branch contains one or more chains of labels $x_1 \leq y_1, \dots, x_m \leq y_m, \dots$; each label that was not already in the endsequent is introduced by a step of $R_{\supset_i}$. By inspection of the rules of **G3I_t**, it is clear that all the formulas that occur in the branch are subformulas of Γ, Δ or formulas of the form $A \supset (B \supset A)$ for some subformula $B \supset A$ of Γ, Δ . To ensure that all proper chains of labels in the reduction tree are finite, it is therefore enough to prove that rule $R_{\supset_i}$ need not be applied twice to the same formula along a chain of labels.

Suppose that we have a chain $x_0 \leq x_1, \dots, x_{n-1} \leq x_n$ in the antecedent and $x_0: A \supset B, x_n: A \supset B$ in the succedent of a branch in the proof search and that $R_{\supset_i}$ has been applied to $x_0: A \supset B$. We need to show that there is no need to apply $R_{\supset_i}$ to $x_n: A \supset B$. Suppose for simplicity that we have a chain of length 2, with $x_0 \equiv x, x_1 \equiv y, x_2 \equiv z$:

$$\frac{\begin{array}{c} x \leq y, y \leq z, y: B \supset (A \supset B), \Gamma' \rightarrow \Delta', z: A \supset B, \\ \vdots \\ x \leq y, y: B \supset (A \supset B), y: A, \Gamma \rightarrow \Delta, y: B \end{array}}{\Gamma \rightarrow \Delta, x: A \supset B} R_{\supset_i}$$

and assume that the top-sequent is closed under all the available rules (excluding $R_{\supset_i}$) of the reduction procedure. We observe that in the application of $L_{\supset}$ on $y: B \supset (A \supset B)$ and $y \leq z$, the right premiss with $z: A \supset B$ both on the left and right is derivable, therefore we only consider the left premiss with $z: B$ is in the succedent. So we have that $y: A$ (as marked) is in the antecedent from the first step of $R_{\supset_i}$ below, and Δ' contains $z: B$ is in the succedent from the application of $L_{\supset}$ on $y: B \supset (A \supset B)$ and $y \leq z$ (we consider only the left premiss since the right premiss with $z: A \supset B$ both on the left and right is derivable). So we can apply Lemma 2.6 and discard $z: A \supset B$.

We can conclude that all the chains of labels in the tree are finite. To conclude that the branch is finite, it is enough to observe that it contains only a finite number of such chains (the number of chains is bounded by a function of the number of disjunctions or commas in the positive part of the endsequent). The general case, where the chain is longer than just $x \leq y, y \leq z$, is similar.

A branch which either ends in an initial sequent or in a sequent with the same labelled formula, even compound, in both the antecedent and succedent, or at the conclusion of $L_{\perp}$, or has a top-sequent amenable to any of the reduction steps, is called *unsaturated*. Every other branch is said to be *saturated*.

2. Construction of the countermodel: If the reduction tree for $\Gamma \rightarrow \Delta$ is not a derivation, it has at least one saturated branch. Let $\Gamma^* \rightarrow \Delta^*$ be the union (respectively, of the antecedents and succedents) of all the sequents $\Gamma_i \rightarrow \Delta_i$ of the branch up to its top-sequent. We define a Kripke model that forces all the formulas in Γ^* and no formula in Δ^* and is therefore a countermodel to the sequent $\Gamma \rightarrow \Delta$.

Consider the frame X , the nodes of which are the labels that appear in the relational atoms in Γ^* and the order on which is given by these relational atoms. Clearly, the construction of the reduction tree imposes the frame properties on the countermodel: $\text{Ref}_{\leq}$ and $\text{Trans}_{\leq}$ hold because the branch is saturated. Moreover, any label that appears in the sequent will appear in a relational atom (and thus in the frame X), because the rule $\text{Ref}_{\leq}$ has been applied. Noetherianity clearly holds because all the strictly ascending chains in the countermodel are finite by construction.

On the frame $(X, \leq)$ we define the following valuation: for each labelled atomic formula $x : P$ in Γ^* we stipulate that $x \Vdash P$. Since the top-sequent is not initial, for all labelled atomic formulas $y : Q$ in Δ^* we infer that $y \nVdash Q$. We then show by induction on $\text{size}(A)$ that $x \Vdash A$ if $x : A$ is in Γ^* and that $x \nVdash A$ if $x : A$ is in Δ^* . Therefore we have a countermodel to the endsequent $\Gamma \rightarrow \Delta$.

- If A is atomic, then the claim holds by the definition of the model.
- If $A \equiv \perp$, it cannot be in Γ^* , by definition of saturated branch: so $x \nVdash A$.
- If $A \equiv B \wedge C$ is in Γ^* , then by the saturation of the branch we also have $x : B$ and $x : C$ in Γ^* . By the induction hypothesis, $x \Vdash B$ and $x \Vdash C$, and therefore $x \Vdash B \wedge C$.
- If $A \equiv B \wedge C$ is in Δ^* , then by the saturation of the branch either $x : B$ or $x : C$ in Δ^* . By the induction hypothesis, $x \nVdash B$ or $x \nVdash C$, and therefore $x \nVdash B \wedge C$.
- If $A \equiv B \vee C$ is in Γ^* , then by the saturation of the branch either $x : B$ or $x : C$ in Γ^* . By the induction hypothesis, $x \Vdash B$ or $x \Vdash C$, and therefore $x \Vdash B \vee C$.
- If $A \equiv B \vee C$ is in Δ^* , then by the saturation of the branch we also have $x : B$ and $x : C$ in Δ^* . By the induction hypothesis, $x \nVdash B$ and $x \nVdash C$, and therefore $x \nVdash B \vee C$.
- If $A \equiv B \supset C$ is in Γ^* , then for any occurrence of $x \leq y$ in Γ^* we find, by saturation and by the construction of the reduction tree, either an occurrence of $y : B$ in Δ^* or an occurrence of $y : C$ in Γ^* . By the induction hypothesis, in the former case $y \nVdash B$, and in the latter $y \Vdash C$, so in both cases $x \Vdash B \supset C$.
- If $A \equiv B \supset C$ is in Δ^* , we consider the step where it is analysed. If $x : C$ is in the succedent of that step (or any succedent below it), then by the induction hypothesis $x \Vdash B$. Since $x \leq x$ is also in Γ^* by construction of the reduction tree, it follows that $x \Vdash B \supset C$. Otherwise there is $x \leq y$ in Γ^* and $y : C$ in Δ^* . By the induction hypothesis $y \nVdash C$, and therefore $x \nVdash A$. $\dashv$

COROLLARY 3.6. *If a sequent $\Gamma \rightarrow \Delta$ is valid in every reflexive, transitive, and Noetherian frame, then it is derivable in $\mathbf{G3I}_t$.*

COROLLARY 3.7. *A formula A is provable in $\mathbf{Int}$ if and only if the sequent $\rightarrow x : A$ is derivable in $\mathbf{G3I}_t$ for some (or any) label x .*

COROLLARY 3.8. *The rule of cut,*

$$\frac{\Gamma \rightarrow \Delta, x : A \quad x : A, \Gamma' \rightarrow \Delta'}{\Gamma, \Gamma' \rightarrow \Delta, \Delta'} \text{Cut}$$

is admissible in $\mathbf{G3I}_t$.

The proof of Theorem 3.5 is also of interest because it establishes the finite model property for $\mathbf{Int}$ and gives a constructive decision procedure for it, i.e., an algorithm that, given a sequent, constructs either a derivation or a countermodel.

$$((P \supset Q) \supset P) \supset P.$$
$$\begin{array}{c}
\vdots \\
\hline
y \leq w, z \leq w, y \leq z, y \leq y, x \leq y, w : P, z : P, y : (P \supset Q) \supset P \rightarrow y : P, z : Q, w : Q \quad \text{L}\supset \\
\text{Trans} \\
\hline
z \leq w, y \leq z, y \leq y, x \leq y, w : P, z : P, y : (P \supset Q) \supset P \rightarrow y : P, z : Q, w : Q \\
\hline
y \leq z, y \leq y, x \leq y, z : P, y : (P \supset Q) \supset P \rightarrow y : P, z : Q, z : P \supset Q \quad \text{R}\supset \\
\hline
\vdots \\
y \leq z, y \leq y, x \leq y, z : P, y : (P \supset Q) \supset P \rightarrow y : P, z : Q \quad \text{L}\supset \\
\hline
\vdots \\
y \leq y, x \leq y, y : (P \supset Q) \supset P \rightarrow y : P, y : P \supset Q \quad \text{R}\supset \\
\hline
\vdots \\
y \leq y, x \leq y, y : (P \supset Q) \supset P \rightarrow y : P \quad \text{L}\supset \\
\hline
\vdots \\
x \leq y, y : (P \supset Q) \supset P \rightarrow y : P \quad \text{Ref}_{\leq} \\
\hline
\rightarrow x : ((P \supset Q) \supset P) \supset P \quad \text{L}\supset
\end{array}$$
$$\begin{array}{c}
\frac{z \leq y, z: Q \supset (P \supset Q), z: P, y \leq y, x \leq y, y: P \supset (((P \supset Q) \supset P) \supset P), y: (P \supset Q) \supset P \rightarrow y: P, z: Q, z: P \supset Q}{z \leq y, z: Q \supset (P \supset Q), z: P, y \leq y, x \leq y, y: P \supset (((P \supset Q) \supset P) \supset P), y: (P \supset Q) \supset P \rightarrow y: P, z: Q} \text{R}_{\supset} \\
\frac{y \leq y, x \leq y, y: P \supset (((P \supset Q) \supset P) \supset P), y: (P \supset Q) \supset P \rightarrow y: P, y: P \supset Q}{y \leq y, x \leq y, y: P \supset (((P \supset Q) \supset P) \supset P), y: (P \supset Q) \supset P \rightarrow y: P, y: (P \supset Q) \supset P \rightarrow y: P} \text{L}_{\supset} \\
\frac{x \leq y, y: P \supset (((P \supset Q) \supset P) \supset P), y: (P \supset Q) \supset P \rightarrow y: P}{\rightarrow x: ((P \supset Q) \supset P) \supset P} \text{Ref}_{\leq} \\
\rightarrow x: ((P \supset Q) \supset P) \supset P \text{R}_{\supset}
\end{array}$$
$$\forall x_1 \geq x (\forall x_2 \geq x_1 (\forall x_3 \geq x_2 (x_3 \Vdash P \Rightarrow x_3 \Vdash Q) \Rightarrow x_2 \Vdash P) \Rightarrow x_1 \Vdash P)$$
$$\forall x_2 \geq y (\forall x_3 \geq x_2 (x_3 \Vdash P \Rightarrow x_3 \Vdash \underline{Q}) \Rightarrow x_2 \Vdash P).$$

§4. Embedding into Grzegorczyk logic. We recall that modal logic is obtained by adding the modal operator $\Box$ to the language of propositional logic, and the inductive clauses for valuations of modal formulas are the following:

Initial sequent

$$x: P, \Gamma \rightarrow \Delta, x: P$$

Propositional rules

$$\frac{x: A, x: B, \Gamma \rightarrow \Delta}{x: A \wedge B, \Gamma \rightarrow \Delta} L\wedge$$

$$\frac{\Gamma \rightarrow \Delta, x: A \quad \Gamma \rightarrow \Delta, x: B}{\Gamma \rightarrow \Delta, x: A \wedge B} R\wedge$$

$$\frac{x: A, \Gamma \rightarrow \Delta \quad x: B, \Gamma \rightarrow \Delta}{x: A \vee B, \Gamma \rightarrow \Delta} L\vee$$

$$\frac{\Gamma \rightarrow \Delta, x: A, x: B}{\Gamma \rightarrow \Delta, x: A \vee B} R\vee$$

$$\frac{\Gamma \rightarrow \Delta, x: A \quad x: B, \Gamma \rightarrow \Delta}{x: A \supset B, \Gamma \rightarrow \Delta} L\supset$$

$$\frac{x: A, \Gamma \rightarrow \Delta, x: B}{\Gamma \rightarrow \Delta, x: A \supset B} R\supset$$

$$\frac{}{x: \perp, \Gamma \rightarrow \Delta} L\perp$$

Modal rules

$$\frac{x \leq y, y: A, x: \Box A, \Gamma \rightarrow \Delta}{x \leq y, x: \Box A, \Gamma \rightarrow \Delta} L\Box$$

$$\frac{x \leq y, y: G(A), \Gamma \rightarrow \Delta, y: A}{\Gamma \rightarrow \Delta, x: \Box A} R\Box Z$$

Mathematical rules

$$\frac{x \leq x, \Gamma \rightarrow \Delta}{\Gamma \rightarrow \Delta} \text{Ref}_{\leq}$$

$$\frac{x \leq z, x \leq y, y \leq z, \Gamma \rightarrow \Delta}{x \leq y, y \leq z, \Gamma \rightarrow \Delta} \text{Trans}_{\leq}$$

TABLE 3. The sequent calculus **G3Grz**. Rule $R\Box Z$ has the condition that y is fresh.

$$x \not\vdash \perp.$$

$$x \vdash A \supset B \text{ if and only if } x \vdash A \Rightarrow x \vdash B.$$

$$x \vdash A \wedge B \text{ if and only if } x \vdash A \text{ and } x \vdash B.$$

$$x \vdash A \vee B \text{ if and only if } x \vdash A \text{ or } x \vdash B.$$

$$x \vdash \Box A \text{ if and only if } \forall y (x \leq y \Rightarrow y \vdash A).$$

The provability logic **Grz** (Grzegorzczuk logic) [1, 6, 10] is an extension of basic modal logic **K** with the additional schemata

$$\Box A \supset A, \quad (\text{Ax. T})$$

$$\Box A \supset \Box \Box A, \quad (\text{Ax. 4})$$

$$\Box(G(A) \supset A) \supset A, \quad (\text{Ax. Grz})$$

where $G(A) \equiv \Box(A \supset \Box A)$. **Grz** is characterised by reflexive, transitive, and Noetherian frames [6]. The sequent calculus **G3Grz** for **Grz** (see Table 3) satisfies all usual structural rules, including hp-invertibility of its rules [6].

As shown in [6], an indirect decision procedure for **Int** is obtained through faithfulness of the embedding of **Int** into **Grz** via the translation $\cdot^\Box$ inductively defined as

$$P^\Box \equiv \Box P,$$

$$\perp^\Box \equiv \perp,$$

$$(A \wedge B)^\Box \equiv A^\Box \wedge B^\Box,$$

$$(A \vee B)^\square \equiv A^\square \vee B^\square,$$

$$(A \supset B)^\square \equiv \square(A^\square \supset B^\square).$$

REMARK 4.1. The translation of $R\supset_t$ is the following:

$$\frac{x \leq y, y: \square(B^\square \supset \square(A^\square \supset B^\square)), y: A^\square, \Gamma^\square \rightarrow \Delta^\square, y: B^\square}{\Gamma^\square \rightarrow \Delta^\square, x: \square(A^\square \supset B^\square)} \quad (y \text{ fresh}).$$

If we set $A \equiv \top$, this is equivalent to

$$\frac{x \leq y, y: \square(B^\square \supset \square B^\square), \Gamma^\square \rightarrow \Delta^\square, y: B^\square}{\Gamma^\square \rightarrow \Delta^\square, x: \square B^\square} \quad (y \text{ fresh})$$

which is an instance of $R\square Z$, the rule that allows decidability in the calculus **G3Grz** for Grzegorczyk logic.

We now want to give a proof of faithfulness alternative to the one is given in [6] by using **G3I_t** in place of **G3I**. We first need a few lemmata:

LEMMA 4.2. *If there is a derivation in **G3Grz** of height n of*

$$x: A \supset B, \Gamma \rightarrow \Delta, \quad (7)$$

then there are derivations of height at most n of

$$\Gamma \rightarrow \Delta, x: A \quad (8)$$

$$x: B, \Gamma \rightarrow \Delta. \quad (9)$$

If, moreover, $x: A \supset B$ is used as the principal formula somewhere in the given derivation of (7), then the derivations of (8) and (9) have height at most $n - 1$.

PROOF. We slightly modify the usual argument for hp-invertibility of $L\supset$ (see, e.g., [13, Proposition 4.11]). The proof proceeds by induction on n .

$n = 0$: Trivial.

$n > 0$: If $x: A \supset B$ is principal in the last rule applied in the derivation of (7), then the two branches are derivations of (8) and (9) of height at most $n - 1$. If it is not principal and the last rule applied is *rule*, then we proceed as usual by applying the induction hypothesis to the previous step(s) followed by *rule*. $\dashv$

LEMMA 4.3. *The rule*

$$\frac{x \leq y, x: A^\square, y: A^\square, \Gamma \rightarrow \Delta}{x \leq y, x: A^\square, \Gamma \rightarrow \Delta}$$

*with the condition that the top-sequent is saturated under transitivity, is hp-admissible in **G3Grz**.*

PROOF. We prove it by induction on the height of the derivation of the premiss, with a subinduction on the length of A .

$n = 0$: Trivial.

$n > 0$: The only nontrivial cases are those in which the last rule applied is a left rule and $y: A^\square$ is principal. Cases $L\wedge$ and $L\vee$ are dealt with as in Lemma 2.5, and

$L\Box$ as in [6, Lemma 3.14]. The assumption of saturation under transitivity makes the application of $\text{Trans}_{\leq}$ in [6, Lemma 3.14] unnecessary, thus ensuring height preservation. $\dashv$

LEMMA 4.4. *The rule*

$$\frac{x: A^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta}{x: A^\Box, x: \Box(B \supset C), \Gamma \rightarrow \Delta}$$

with the condition that the top-sequent is saturated under transitivity, is *hp-admissible* in **G3Grz**.

PROOF. Suppose that there is a derivation of

$$x: A^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta \quad (10)$$

of height n . We prove by induction on n that there is a derivation of

$$x: A^\Box, x: \Box(B \supset C), \Gamma \rightarrow \Delta \quad (11)$$

of height n .

$n = 0$: All cases are trivial.

$n > 0$: The cases in which the principal formula is in Γ or Δ are trivial.

Suppose that the principal formula is $x: A^\Box$, and consider the case in which $A \equiv A_1 \wedge A_2$, which means that we have a derivation

$$\frac{x: A_1^\Box, x: A_2^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta}{x: A^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta} L\wedge$$

We can apply hp-weakening to the premiss and get

$$x: A^\Box, x: A_1^\Box, x: A_2^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta,$$

to which we can apply the induction hypothesis:

$$x: A^\Box, x: A_1^\Box, x: A_2^\Box, x: \Box(B \supset C), \Gamma \rightarrow \Delta.$$

We conclude by $L\wedge$ and contraction.

Suppose that the principal formula is $x: A^\Box$, and consider the case in which $A \equiv A_1 \vee A_2$, which means that we have a derivation

$$\frac{x: A_1^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta \quad x: A_2^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta}{x: A^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta} L\vee$$

We can apply hp-weakening to the premisses and get

$$\begin{aligned} x: A^\Box, x: A_1^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta, \\ x: A^\Box, x: A_2^\Box, x: \Box((A^\Box \supset B) \supset C), \Gamma \rightarrow \Delta, \end{aligned}$$

to which we can apply the induction hypothesis:

$$\begin{aligned} x: A^\Box, x: A_1^\Box, x: \Box(B \supset C), \Gamma \rightarrow \Delta, \\ x: A^\Box, x: A_2^\Box, x: \Box(B \supset C), \Gamma \rightarrow \Delta. \end{aligned}$$

We conclude by $L\vee$ and contraction.

Suppose that the principal formula is $x: A^\square$, and consider the case in which $A \equiv A_1 \supset A_2$ or $A \equiv P$, which means that we have a derivation

$$\frac{y: A, x: A^\square, x: \square((A^\square \supset B) \supset C), x \leq y, \Gamma' \rightarrow \Delta}{x: A^\square, x: \square((A^\square \supset B) \supset C), x \leq y, \Gamma' \rightarrow \Delta} \text{L}\square$$

where $\Gamma \equiv x \leq y, \Gamma'$. We can apply the induction hypothesis to the premiss:

$$y: A, x: A^\square, x: \square(B \supset C), x \leq y, \Gamma' \rightarrow \Delta.$$

We conclude by $\text{L}\supset$.

Now suppose that the principal formula is $x: \square((A^\square \supset B) \supset C)$. This means that we have

$$\frac{y: (A^\square \supset B) \supset C, x: A^\square, x: \square((A^\square \supset B) \supset C), x \leq y, \Gamma' \rightarrow \Delta}{x: A^\square, x: \square((A^\square \supset B) \supset C), x \leq y, \Gamma' \rightarrow \Delta} \text{L}\square$$

where $\Gamma \equiv x \leq y, \Gamma'$. We can assume that $y: (A^\square \supset B) \supset C$ is used as the principal formula somewhere above this instance of $\text{L}\square$: if not, then we could find a derivation of (10) without this instance of $\text{L}\square$, this would have smaller height and therefore we could apply the induction hypothesis to it. By applying hp-weakening to the premiss, we obtain a derivation of

$$y: A^\square, y: (A^\square \supset B) \supset C, x: A^\square, x: \square((A^\square \supset B) \supset C), x \leq y, \Gamma' \rightarrow \Delta$$

of height $n - 1$ and such that $y: (A^\square \supset B) \supset C$ is used as the principal formula somewhere above. Now by Lemma 4.2 on invertibility of $\text{L}\supset$ we get derivations of

$$y: C, y: A^\square, x: A^\square, x: \square((A^\square \supset B) \supset C), x \leq y, \Gamma' \rightarrow \Delta, \quad (12)$$

$$y: A^\square, x: A^\square, x: \square((A^\square \supset B) \supset C), x \leq y, \Gamma' \rightarrow \Delta, y: A^\square \supset B, \quad (13)$$

both of height $n - 2$. Now we can apply the induction hypothesis to (12) and get a derivation of

$$y: C, y: A^\square, x: A^\square, x: \square(B \supset C), x \leq y, \Gamma' \rightarrow \Delta \quad (14)$$

of height $n - 2$. By applying hp-invertibility of $\text{R}\supset$ and hp-contraction to (13), we get a derivation of

$$y: A^\square, x: A^\square, x: \square((A^\square \supset B) \supset C), x \leq y, \Gamma' \rightarrow \Delta, y: B$$

of height $n - 2$, to which we can apply the induction hypothesis and get a derivation of

$$y: A^\square, x: A^\square, x: \square(B \supset C), x \leq y, \Gamma' \rightarrow \Delta, y: B \quad (15)$$

of height $n - 2$. Now we can apply $\text{L}\supset$ to (14) and (15) and get a derivation of

$$y: B \supset C, y: A^\square, x: A^\square, x: \square(B \supset C), x \leq y, \Gamma' \rightarrow \Delta$$

of height $n - 1$, which by an application of $\text{L}\square$ gives a derivation of

$$y: A^\square, x: A^\square, x: \square(B \supset C), x \leq y, \Gamma' \rightarrow \Delta$$

of height n . We conclude by Lemma 4.3. $\dashv$

Now we are able to prove faithfulness:

THEOREM 4.5 (Faithfulness). *Let Γ, Δ be multisets of labelled formulas in the language of **G3I_t**, Γ', Δ' multisets of labelled atomic formulas, with Γ' possibly containing also relational atoms. If*

$$\mathbf{G3Grz} \vdash \Gamma^\square, \Gamma' \rightarrow \Delta^\square, \Delta',$$

then

$$\mathbf{G3I_t} \vdash \Gamma, \Gamma' \rightarrow \Delta, \Delta'.$$

PROOF. By induction on the height of the derivation of $\Gamma^\square, \Gamma' \rightarrow \Delta^\square, \Delta'$. We assume that $\Gamma^\square, \Gamma' \rightarrow \Delta^\square, \Delta'$ is saturated under transitivity: this can be done without loss of generality since it is equivalent to apply $\text{Trans}_{\leq}$ in the proof search as soon as possible, which is innocuous because the rule operates on labels already introduced.

$n = 0$: If it is an initial sequent or the conclusion of $\text{L}\perp$, then it can be translated smoothly into the corresponding initial sequent or rule in **G3I_t**.

$n > 0$: First, notice that rules for $\supset$ cannot produce a sequent of this form. If it is the conclusion of a rule for $\perp, \wedge, \vee$, then it can be translated smoothly into the corresponding initial sequent or rule in **G3I_t**. If it is derived by a modal rule, then the principal formula can be of the form $\Box P$ or of the form $\Box(A^\square \supset B^\square)$. We have four cases:

– If $\Box P$ is principal on the left, we have (with $\Gamma = x : P, \Gamma''$)

$$\frac{x \leq y, y : P, x : \Box P, \Gamma''^\square, \Gamma' \rightarrow \Delta^\square, \Delta'}{x \leq y, x : \Box P, \Gamma''^\square, \Gamma' \rightarrow \Delta^\square, \Delta'} \text{L}\Box$$

which, using the induction hypothesis, is translated into the admissible **G3I_t** step

$$\frac{x \leq y, y : P, x : P, \Gamma'', \Gamma' \rightarrow \Delta, \Delta'}{x \leq y, x : P, \Gamma'', \Gamma' \rightarrow \Delta, \Delta'}.$$

– If $\Box P$ is principal on the right, we have (with $\Delta = x : P, \Delta''$)

$$\frac{x \leq y, y : G(P), \Gamma^\square, \Gamma' \rightarrow \Delta''^\square, y : P, \Delta'}{\Gamma^\square, \Gamma' \rightarrow \Delta''^\square, x : \Box P, \Delta'} \text{R}\Box\text{Z}$$

which, as seen in Remark 4.1, is the translation of a step of rule $\text{R}\supset_t$ with $\top \supset P$ as the principal formula.

– If $\Box(A^\square \supset B^\square)$ is principal on the left, we have (with $\Gamma = A \supset B, \Gamma''$ and $\Gamma' = x \leq y, \Gamma'''$)

$$\frac{x \leq y, y : A^\square \supset B^\square, x : \Box(A^\square \supset B^\square), \Gamma''^\square, \Gamma''' \rightarrow \Delta^\square, \Delta'}{x \leq y, x : \Box(A^\square \supset B^\square), \Gamma''^\square, \Gamma''' \rightarrow \Delta^\square, \Delta'} \text{L}\Box$$

from which, by hp-invertibility of $\text{L}\supset$ in **G3Grz** we have

$$\mathbf{G3Grz} \vdash x \leq y, y : B^\square, x : \Box(A^\square \supset B^\square), \Gamma''^\square, \Gamma''' \rightarrow \Delta^\square, \Delta',$$

$$\mathbf{G3Grz} \vdash x \leq y, x : \Box(A^\square \supset B^\square), \Gamma''^\square, \Gamma''' \rightarrow \Delta^\square, y : A^\square, \Delta',$$

to which the induction hypothesis applies:

$$\begin{aligned}\mathbf{G3I}_t &\vdash x \leq y, y: B, x: A \supset B, \Gamma'', \Gamma''' \rightarrow \Delta, \Delta', \\ \mathbf{G3I}_t &\vdash x \leq y, x: A \supset B, \Gamma'', \Gamma''' \rightarrow \Delta, y: A, \Delta'.\end{aligned}$$

We conclude by an application of $L\supset$.

- If $\Box(A^\Box \supset B^\Box)$ is principal on the right, we have (with $\Delta = x: A \supset B, \Delta''$)

$$\frac{x \leq y, y: G(A^\Box \supset B^\Box), \Gamma^\Box, \Gamma' \rightarrow \Delta''^\Box, y: A^\Box \supset B^\Box, \Delta'}{\Gamma^\Box, \Gamma' \rightarrow \Delta''^\Box, x: \Box(A^\Box \supset B^\Box), \Delta'} \text{R}\Box Z$$

from which, by hp-invertibility of $R\supset$ in $\mathbf{G3Grz}$ we have

$$\mathbf{G3Grz} \vdash x \leq y, y: G(A^\Box \supset B^\Box), y: A^\Box, \Gamma^\Box, \Gamma' \rightarrow \Delta''^\Box, y: B^\Box, \Delta'.$$

By Lemma 4.4, it follows that

$$\mathbf{G3Grz} \vdash x \leq y, y: \Box(B^\Box \supset \Box(A^\Box \supset B^\Box)), y: A^\Box, \Gamma''^\Box, \Gamma' \rightarrow \Delta''^\Box, y: B^\Box, \Delta',$$

to which the induction hypothesis applies:

$$\mathbf{G3I}_t \vdash x \leq y, y: B \supset (A \supset B), y: A, \Gamma, \Gamma' \rightarrow \Delta'', y: B, \Delta'.$$

We conclude by $R\supset_t$. ⊢

§5. Future work. Since the logic $\mathbf{Grz}$ studied in the present paper is characterised by reflexive, transitive, and Noetherian frames, we also plan to use the approach of [7] to define a variant of induction principle, which we may dub *Grzegorzcyk induction* corresponding to rule $R\Box Z$:

$$\forall x [\forall y \leq x (GE(y) \implies E(y)) \implies \forall y \leq x E(y)],$$

where $GE(y)$ is an abbreviation for $\forall z \leq y (E(z) \implies \forall w \leq z E(w))$. This can be considered a weak form of induction compatible with reflexivity, and may give a different perspective of the semantics of both $\mathbf{Grz}$ and $\mathbf{Int}$ and may give some insights on the properties of the accessibility relation.

We then plan to extend the approach of this paper to extensions of $\mathbf{Int}$, such as intermediate logics [5, 16], modal intuitionistic logic [12] and possibly bi-intuitionistic logic [18].

Acknowledgements. Both authors are members of the “Gruppo Nazionale per le Strutture Algebriche, Geometriche e le loro Applicazioni” (GNSAGA) of the Istituto Nazionale di Alta Matematica (INdAM). Last but not least, the authors are grateful to Giovanna Corsi, Eugenio Orlandelli, Mario Piazza, Matteo Tesi, and Tarmo Uustalu for their interest and valuable suggestions.

Funding. Support for this research was granted by the projects “Reducing complexity in algebra, logic, combinatorics—REDCOM” belonging to the programme “Ricerca Scientifica di Eccellenza 2018” of the Fondazione Cariverona, “Infinity and Intensionality: Towards A New Synthesis” funded by the Research Council of Norway, and “Modalities in Substructural Logics: Theory, Methods and

Applications – MOSAIC”, funded by the Community Research and Development Information Service (CORDIS) of the European Commission. The research by Sara Negri was supported in part by the MIUR Excellence Department Project awarded to Dipartimento di Matematica, Università di Genova, CUP D33C23001110001.

REFERENCES

- [1] A. AVRON, *On modal systems having arithmetical interpretations*, this JOURNAL, vol. 49 (1984), no. 3, pp. 935–942. 1984 <http://www.jstor.org/stable/2274147>.
- [2] G. CORSI, *The a fortiori rule: The key to reach termination in intuitionistic logic*, *Logic and Philosophy in Italy. Some Trends and Perspectives. Essays in Honor of Corrado Mangione on his 75th Birthday* (E. Ballo and M. Franchella, editors), Polimettrica, Milano, 2006, pp. 27–47.
- [3] G. CORSI and G. TASSI, *Intuitionistic logic freed of all metarules*, this JOURNAL, vol. 72 (2007), no. 4, pp. 1204–1218. 2007 <https://doi.org/10.2178/jsl/1203350782>.
- [4] R. DYCKHOFF, *Intuitionistic decision procedures since gentzen*, *Advances in Proof Theory. Progress in Computer Science and Applied Logic*, vol. 28 (R. Kahle, T. Strahm, and T. Studer, editors), Birkhäuser, Cham, 2016, pp. 245–267.
- [5] R. DYCKHOFF and S. NEGRI, *Proof analysis in intermediate propositional logics*, *Archive for Mathematical Logic*, vol. 51 (2012), nos. 1–2, pp. 71–92.
- [6] ———, *A cut-free sequent system for Grzegorczyk logic, with an application to the Gödel–McKinsey–Tarski embedding*, *Journal of Logic and Computation*, vol. 26 (2016), no. 1, pp. 169–187.
- [7] G. FELLIN, S. NEGRI, and P. SCHUSTER, *Modal logic for induction*, *Advances in Modal Logic*, vol. 13 (N. Olivetti, R. Verbrugge, S. Negri, and G. Sandu, editors), College Publications, London, 2020, pp. 209–227.
- [8] G. GENTZEN, *Untersuchungen über das logische Schließen I. Mathematische Zeitschrift*, vol. 39 (1934), pp. 176–210.
- [9] ———, *Untersuchungen über das logische Schließen II. Mathematische Zeitschrift*, vol. 39 (1934), pp. 405–431.
- [10] A. GRZEGORCZYK, *Some relational systems and the associated topological spaces*, *Fundamenta Mathematicae*, vol. 60 (1967), pp. 223–231.
- [11] S. A. KRIPKE, *Semantical analysis of modal logic. I. Normal modal propositional calculi*, *Zeitschrift für Mathematische Logik und Grundlagen der Mathematik*, vol. 9 (1963), pp. 67–96.
- [12] P. MAFFEZIOLI, A. NAIBO, and S. NEGRI, *The Church–Fitch knowability paradox in the light of structural proof theory*, *Synthese*, vol. 190 (2013), no. 14, pp. 2677–2716.
- [13] S. NEGRI, *Proof analysis in modal logic*, *Journal of Philosophical Logic*, vol. 34 (2005), nos. 5–6, pp. 507–544.
- [14] ———, *Proof analysis in non-classical logics*, *Logic Colloquium 2005*, (C. Dimitracopoulos, L. Newelski, D. Normann, J. R. Steel, editors), Lecture Notes in Logic, 28, Association for Symbolic Logic, Urbana, 2008, pp. 107–128.
- [15] ———, *On the duality of proofs and countermodels in labelled sequent calculi*, *Automated Reasoning with Analytic Tableaux and Related Methods, TABLEUX 2013* (D. Galmiche and D. Larchey-Wendling, editors), Lecture Notes in Computer Science, 8123, Springer, Berlin, 2013, pp. 5–9.
- [16] ———, *Proofs and countermodels in non-classical logics*, *Logica Universalis*, vol. 8 (2014), no. 1, pp. 25–60.
- [17] S. NEGRI and J. VON PLATO, *Proof Analysis. A Contribution to Hilbert’s Last Problem*, Cambridge University Press, Cambridge, 2011.
- [18] L. PINTO and T. UUSTALU, *A proof-theoretic study of bi-intuitionistic propositional sequent calculus*, *Journal of Logic and Computation*, vol. 28 (2018), no. 1, pp. 165–202. <https://doi.org/10.1093/logcom/exx044>.
- [19] J. VON PLATO, *Saved from the Cellar: Gerhard Gentzen’s Shorthand Notes on Logic and Foundations of Mathematics*, Studies and Sources in the History of Mathematics and Physical Sciences, Springer, London, 2017. <https://doi.org/10.1007/978-3-319-42120-9>.
- [20] G. TAKEUTI, *Proof Theory*, second ed., Studies in Logic and the Foundations of Mathematics, 81, North-Holland, Amsterdam, 1987. With an appendix containing contributions by Georg Kreisel, Wolfram Pohlers, Stephen G. Simpson and Solomon Feferman.

DIPARTIMENTO DI INGEGNERIA
UNIVERSITÀ DI BRESCIA
DELL'INFORMAZIONE
VIA BRANZE 38 - 25123 BRESCIA

and

DIPARTIMENTO DI INFORMATICA
UNIVERSITÀ DI VERONA
STRADA LE GRAZIE 15, 37134 VERONA, ITALY

E-mail: giulio.fellin@univr.it

E-mail: giulio.fellin@unibs.it

DIPARTIMENTO DI MATEMATICA
DIPARTIMENTO DI ECCELLENZA 2023–2027
UNIVERSITÀ DI GENOVA
VIA DODECANESO, 35, 16146 GENOVA, ITALY

E-mail: sara.negri@unige.it